

MC4 Receptor Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54493

Specification

MC4 Receptor Polyclonal Antibody - Product Information

Application WB, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat
Host
Clonality
Calculated MW
Physical State
R32245
Rat
Rabbit
Polyclonal
Rabbit
Polyclonal
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human MC4 Receptor

Epitope Specificity 1-50/332 Isotype IgG

Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane; Multi-pass membrane

protein.

SIMILARITY Belongs to the G-protein coupled receptor

1 family.

SUBUNIT Interacts with ATRNL1. Interacts with

MGRN1, but does not undergo

MGRN1-mediated ubiquitination; this interaction competes with GNAS-binding and thus inhibits agonist-induced cAMP

production.

DISEASE Defects in MC4R are a cause of obesity

(OBESITY) [MIM:601665]. It is a condition characterized by an increase of body weight beyond the limitation of skeletal and physical requirements, as the result of

excessive accumulation of body fat.
This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Important Note

Background Descriptions

The melanocortin family comprises the Alpha-, Beta- and Gamma- melanocyte stimulating hormones (MSH) and adrenocorticotrophin. The receptors for these hormones are seven-transmembrane G protein-coupled proteins that activate adenylyl cyclase. Five melanocortin receptors have been cloned and shown to exhibit different affinities and patterns of expression. MC1-R (MSH-R) is expressed in melanocytes and corticoadrenal tissue. MC2-R is the ACTH receptor and is expressed primarily in the adrenal cortex. MC3-R has been found in specific regions of the brain and is also expressed in placenta and gut. MC4-R is expressed primarily in brain, while MC5-R is expressed at low levels in most tissues.



MC4 Receptor Polyclonal Antibody - Additional Information

Gene ID 4160

Other Names

Melanocortin receptor 4, MC4-R, MC4R

Target/Specificity

Brain, placental, and gut tissues.

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MC4 Receptor Polyclonal Antibody - Protein Information

Name MC4R

Function

Hormone receptor that acts as a key component of the leptin- melanocortin pathway at the intersection of homeostatic maintenance of energetic state (PubMed:32327598, PubMed:33858992). Plays a role in regulating food intake: activation by a stimulating hormone such as anorexigenic alpha-melanocyte stimulating hormone (alpha-MSH) inhibits appetite, whereas binding to a natural antagonist like Agouti-related protein/AGRP promotes appetite. G-protein-coupled receptor that activates conventional Galphas signaling leading to induction of anorexogenic signaling in the hypothalamus to result in negative energy balance (PubMed: 33858992). Regulates the firing activity of neurons from the hypothalamus by alpha-MSH and AGRP independently of Galphas signaling by ligand-induced coupling of closure of inwardly rectifying potassium channel KCNI13 (By similarity). In intestinal epithelial cells, plays a role in the inhibition of hepatic glucose production via nesfatin-1/NUCB2 leading to increased cyclic adenosine monophosphate (cAMP) levels and glucagon-like peptide 1 (GLP-1) secretion in the intestinal epithelium (PubMed: 39562740).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Brain, placental, and gut tissues.


MC4 Receptor Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.



- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MC4 Receptor Polyclonal Antibody - Images




Sample:

Embryo (Mouse) Lysate at 40 ug

Primary: Anti- MC4 Receptor (bs-11417R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 37 kD Observed band size: 52 kD




Sample: MCF-7 Cell (Human) Lysate at 40 ug

Primary: Anti-MC4 Receptor (bs-11417R) at 1/300 dilution




Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 37 kD Observed band size: 40 kD



Positive control: RSC96

Isotype Control Antibody: Rabbit IgG ; Secondary Antibody: Goat anti-rabbit IgG-FITC, Dilution: 1:100 in 1 X PBS containing 0.5% BSA ; Primary Antibody Dilution: 6 μ g in 100 μ L1X PBS containing 0.5% BSA.




Blank control: 293T(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice).

Primary Antibody: Rabbit Anti-MC4 Receptor /AF488 Conjugated antibody (bs-11417R-AF488), Dilution: 1 μ g in 100 μ L 1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG/AF488(orange) ,used under the same conditions.





Blank control: 293T(fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice). Primary Antibody: Rabbit Anti-MC4 Receptor /AF488 Conjugated antibody (bs-11417R-AF488), Dilution: 1 μ g in 100 μ L 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG/AF488(orange) ,used under the same conditions.