

Dscam Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54510**Specification**

Dscam Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O60469
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	222260

Dscam Polyclonal Antibody - Additional Information**Gene ID** 1826**Other Names**

Down syndrome cell adhesion molecule, CHD2, DSCAM

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Dscam Polyclonal Antibody - Protein Information**Name** DSCAM**Function**

Cell adhesion molecule that plays a role in neuronal self- avoidance. Promotes repulsion between specific neuronal processes of either the same cell or the same subtype of cells. Mediates within retinal amacrine and ganglion cell subtypes both isoneuronal self- avoidance for creating an orderly dendritic arborization and heteroneuronal self-avoidance to maintain the mosaic spacing between amacrine and ganglion cell bodies (PubMed:10925149). Receptor for netrin required for axon guidance independently of and in collaboration with the receptor DCC. Might also collaborate with UNC5C in NTN1- mediated axon repulsion independently of DCC (By similarity). In spinal cord development plays a role in guiding commissural axons projection and pathfinding across the ventral midline to reach the floor plate upon ligand binding (PubMed:18585357, PubMed:19196994). Mediates intracellular signaling by stimulating the activation of MAPK8 and MAP kinase p38 (PubMed:18585357, PubMed:19196994). Adhesion molecule that promotes lamina-specific synaptic connections in the retina: expressed in specific

subsets of interneurons and retinal ganglion cells (RGCs) and promotes synaptic connectivity via homophilic interactions (By similarity).

Cellular Location

[Isoform Short]: Secreted.


Tissue Location

Primarily expressed in brain.


Dscam Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dscam Polyclonal Antibody - Images

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Dscam) Polyclonal Antibody, Unconjugated (bs-11456R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Dscam) Polyclonal Antibody, Unconjugated (bs-11456R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.