

SEMA4F Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54518**Specification**

SEMA4F Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O95754
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SEMA4F
Epitope Specificity	239-288/770
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein
SIMILARITY	Belongs to the semaphorin family. Contains 1 Ig-like C2-type (immunoglobulin-like) domain. Contains 1 PSI domain. Contains 1 Sema domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Semaphorins are a family of cell surface and secreted proteins that are conserved from insects to humans. Members of this family of proteins are approximately 750 amino acids in length (including signal sequences) and are defined by a conserved extracellular "semaphorin" domain of approximately 500 amino acids containing 14-16 cysteines, blocks of conserved sequences and no obvious repeats. Secreted and cell-bound semaphorins chemically attract and repel the growth of neural axons, guiding the development of intricate networks of neural tissue. SEMA4F (semaphorin-4F), also known as SEMAM, SEMAW or PRO2353, is a 770 amino acid member of the semaphorin family. Localized to the membrane, SEMA4F is a single-pass type I protein that is involved in growth cone collapse of retinal ganglion-cell axons. SEMA4F is highly expressed in postnatal brain and lung and contains one immunoglobulin-like (Ig-like) domain, one PSI domain and one semaphorin domain. Two isoforms exist due to alternative splicing events.

SEMA4F Polyclonal Antibody - Additional Information**Gene ID** 10505**Other Names**

Semaphorin-4F, Semaphorin-M, Sema M, Semaphorin-W, Sema W, SEMA4F, SEMAM, SEMAW

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SEMA4F Polyclonal Antibody - Protein Information

Name SEMA4F

Synonyms SEMAM, SEMAW

Function

Probable cell surface receptor that regulates oligodendroglial precursor cell migration (By similarity). Might also regulate differentiation of oligodendroglial precursor cells (By similarity). Has growth cone collapse activity against retinal ganglion-cell axons (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Postsynaptic density {ECO:0000250|UniProtKB:Q9Z143}. Perikaryon {ECO:0000250|UniProtKB:Q9Z123}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9Z123}. Note=Colocalizes with DLG4 at synapses {ECO:0000250|UniProtKB:Q9Z123}

SEMA4F Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SEMA4F Polyclonal Antibody - Images