

VAX2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54523

Specification

VAX2 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9UIW0
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	31 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human VAX2
Epitope Specificity	131-205/290
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nuclear
SIMILARITY	Belongs to the EMX homeobox family. Contains 1 homeobox DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

VAX2 is a 290 amino acid homeobox protein that plays a crucial role in development of the eye, particularly in the specification of the ventral optic vesicle and in establishment of a correct dorsoventral pattern. VAX2 acts as a transcription factor with VAX1 to cooperatively regulate retinal differentiation, neuroepithelial cell proliferation and axial polarization in the retina. Together, VAX1 and VAX2 repress transcription of Pax-6, a strong inducer of retinal development. VAX1 and VAX2 localize to the nucleus and contain one homeobox DNA-binding domain each.

VAX2 Polyclonal Antibody - Additional Information

Gene ID 25806

Other Names

Ventral anterior homeobox 2, VAX2

Dilution

WB~~1:1000
IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
ICC~~N/A

\>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

VAX2 Polyclonal Antibody - Protein Information

Name VAX2

Function

Transcription factor that may function in dorsoventral specification of the forebrain. Regulates the expression of Wnt signaling antagonists including the expression of a truncated TCF7L2 isoform that cannot bind CTNNB1 and acts therefore as a potent dominant-negative Wnt antagonist. Plays a crucial role in eye development and, in particular, in the specification of the ventral optic vesicle (By similarity). May be a regulator of axial polarization in the retina.

Cellular Location

Nucleus.

VAX2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VAX2 Polyclonal Antibody - Images