

**NHLH2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54532****Specification****NHLH2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q02577</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human NHLH2
Epitope Specificity	1-50/135
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus
SIMILARITY	Contains 1 bHLH (basic helix-loop-helix) domain.
SUBUNIT	Efficient DNA binding requires dimerization with another bHLH protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

The helix-loop-helix (HLH) structures are known motifs commonly found in membrane-active and DNA-binding proteins. The helix-loop-helix proteins HEN1 and HEN2 are DNA-binding proteins that may be involved in cell-type determination in the early nervous system. Studies of expression in normal tissues have demonstrated expression of NHLH1/NSCL-1 and NHLH2/NSCL-2, the genes encoding HEN1 and HEN2, in the developing central and peripheral nervous system, specifically in developing neurons.

**NHLH2 Polyclonal Antibody - Additional Information****Gene ID** 4808**Other Names**

Helix-loop-helix protein 2, HEN-2, Class A basic helix-loop-helix protein 34, bHLHa34, Nescient helix loop helix 2, NSCL-2, NHLH2, BHLHA34, HEN2, KIAA0490

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class

=<span class="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class="dilution\_IF">IF~~1:50~200</span><br \><span class="dilution\_ICC">ICC~~N/A</span><br \><span class="dilution\_E">E~~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**NHLH2 Polyclonal Antibody - Protein Information**

**Name** NHLH2

**Synonyms** BHLHA34, HEN2, KIAA0490

**Function**

Transcription factor which binds the E box motif 5'- CA[TC][AG]TG-3'. Involved in regulating energy expenditure, body mass, voluntary physical activity, mating behavior and reproductive longevity, acting through the hypothalamic-pituitary-gonadal axis. Acts as a transcriptional activator of target genes, including NDN, PCSK1, MC4R (By similarity). Is also a transcriptional activator of KISS1 (PubMed:<a href="http://www.uniprot.org/citations/35066646" target="\_blank">35066646</a>). May act centrally to regulate function of both white and brown adipose tissue. Together with NHLH1, required to maintain migration and survival of cells in the anterior extramural migration stream (aes), which forms the precerebellar nuclei. Also, in concert with NHLH1, may determine fate of gonadotropin releasing hormone-1 (GnRH-1) neurons.


**Cellular Location**


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.

**NHLH2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**NHLH2 Polyclonal Antibody - Images**



Sample:

PC-3 (human) cell Lysate at 40 ug

Lovo (human) cell Lysate at 40 ug

Primary: Anti- NHLH2 (bs-11524R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 15 kD

Observed band size: 18 kD