

Alpha chimerin Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) **Catalog # AP54538**

Specification

Alpha chimerin Polyclonal Antibody - Product Information

Application **Primary Accession**

Reactivity Host Clonality Calculated MW Physical State Immunogen

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

WB, IHC-P, IHC-F, IF, ICC, E

P15882

Rat, Dog, Bovine

Rabbit **Polyclonal 53 KDa** Liquid

KLH conjugated synthetic peptide derived

from human Alpha-chimerin

151-250/459

laG

0.01M TBS (pH7.4) with 1% BSA, 0.02% Buffer

Proclin300 and 50% Glycerol.

SIMILARITY Contains 1 phorbol-ester/DAG-type zinc

finger. Contains 1 Rho-GAP domain.

Contains 1 SH2 domain.

SUBUNIT Interacts with EPHA4: effector of EPHA4 in

axon guidance linking EPHA4 activation to

RAC1 regulation

DISEASE Defects in CHN1 are the cause of Duane

retraction syndrome type 2 (DURS2)

[MIM:604356]. Duane retraction syndrome is a congenital eye movement disorder characterized by a failure of cranial nerve

VI (the abducens nerve) to develop normally, resulting in restriction or

absence of abduction, adduction, or both, and narrowing of the palpebral fissure and

retraction of the globe on attempted adduction. Undiagnosed in children, it can

lead to amblyopia, a permanent uncorrectable loss of vision.

This product as supplied is intended for Important Note research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

This gene encodes GTPase-activating protein for ras-related p21-rac and a phorbol ester receptor. It is predominantly expressed in neurons, and plays an important role in neuronal signal-transduction mechanisms. Mutations in this gene are associated with Duane's retraction syndrome 2 (DURS2). Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeg, Apr 2011]



Alpha chimerin Polyclonal Antibody - Additional Information

Gene ID 1123

Other Names

N-chimaerin, A-chimaerin, Alpha-chimerin, N-chimerin, NC, Rho GTPase-activating protein 2, CHN1, ARHGAP2, CHN

Target/Specificity

In neurons in brain regions that are involved in learning and memory processes.

Dilution

```
<span class ="dilution_WB">WB~~1:1000</span><br \><span class
="dilution_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution_IF">IF~~1:50~200</span><br \><span class ="dilution_ICC">ICC~~N/A</span><br \><span class ="dilution_E">E~~N/A</span>
```

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Alpha chimerin Polyclonal Antibody - Protein Information

Name CHN1

Synonyms ARHGAP2, CHN

Function

GTPase-activating protein for p21-rac and a phorbol ester receptor. Involved in the assembly of neuronal locomotor circuits as a direct effector of EPHA4 in axon guidance.

Tissue Location

In neurons in brain regions that are involved in learning and memory processes

Alpha chimerin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Alpha chimerin Polyclonal Antibody - Images



