

FAM126A Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54542

Specification

FAM126A Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession
Reactivity Rat, Pig, Dog
Host Rabbit
Clonality Polyclonal
Calculated MW 57625

FAM126A Polyclonal Antibody - Additional Information

Gene ID 84668

Other Names

Hyccin, Down-regulated by CTNNB1 protein A, Protein FAM126A, FAM126A, DRCTNNB1A {ECO:0000303|PubMed:10910037}

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FAM126A Polyclonal Antibody - Protein Information

Name HYCC1 (HGNC:24587)

Function

Component of a complex required to localize phosphatidylinositol 4-kinase (PI4K) to the plasma membrane (PubMed:26571211). The complex acts as a regulator of phosphatidylinositol 4-phosphate (PtdIns(4)P) synthesis (PubMed:26571211). HYCC1 plays a key role in oligodendrocytes formation, a cell type with expanded plasma membrane that requires generation of PtdIns(4)P (PubMed:26571211). Its role in oligodendrocytes formation probably explains its importance in myelination of the central and



peripheral nervous system (PubMed:16951682, PubMed:26571211). May also have a role in the beta- catenin/Lef signaling pathway (Probable).

Cellular Location

Cytoplasm, cytosol. Cell membrane Note=Localizes to the cytosol and is recruited to the plasma membrane following interaction with other components of the phosphatidylinositol 4-kinase (PI4K) complex.

Tissue Location


Widely expressed. Highest levels in heart, brain, placenta, spleen and testis.

FAM126A Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

FAM126A Polyclonal Antibody - Images



Sample:

Heart (Mouse) Lysate at 40 ug

Primary: Anti-FAM126A (bs-11554R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 58 kD Observed band size: 58 kD