

RND2/Rho7 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54556**Specification**

RND2/Rho7 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	P52198
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25369

RND2/Rho7 Polyclonal Antibody - Additional Information**Gene ID** 8153**Other Names**

Rho-related GTP-binding protein RhoN, Rho family GTPase 2, Rho-related GTP-binding protein Rho7, Rnd2, RND2, ARHN, RHO7

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RND2/Rho7 Polyclonal Antibody - Protein Information**Name** RND2**Synonyms** ARHN, RHO7**Function**

May be specifically involved in neuronal and hepatic functions. Is a C3 toxin-insensitive member of the Rho subfamily (By similarity).

Cellular Location

Cytoplasmic vesicle, secretory vesicle, acrosome membrane; Lipid-anchor; Cytoplasmic side.
Note=Colocalizes with RACGAP1 in Golgi-derived proacrosomal vesicles and the acrosome.


Tissue Location

Highly expressed in testis.

RND2/Rho7 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RND2/Rho7 Polyclonal Antibody - Images


Sample: Brain (Mouse) Lysate at 40 ug

Primary: Anti-RND2/Rho7 (bs-11583R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 25 kD

Observed band size: 26 kD



Sample: Cerebellum (Mouse) Lysate at 40 ug
Primary: Anti-RND2/Rho7 (bs-11583R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 25 kD
Observed band size: 26 kD