

**TULP3 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54565****Specification****TULP3 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">075386</a>
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TULP3
Epitope Specificity	251-350/442
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus. Cell membrane. Cell projection, cilium (By similarity). Cytoplasm (By similarity). Secreted (By similarity). Note=Does not have a cleavable signal peptide and is secreted by a non-conventional pathway (By similarity). Translocates from the plasma membrane to the nucleus upon activation of guanine nucleotide-binding protein G(q) subunit alpha.
SIMILARITY	Belongs to the TUB family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

Mutations in the mouse Tub gene gradually lead to obesity, strongly resembling the late-onset obesity observed in the human population. In addition to excessive deposition of adipose tissue, mice with the Tub phenotype also suffer retinal degeneration and neurosensory hearing loss. A human homolog of the Tub gene has been identified, as have three related proteins, called Tubby-like protein 1 (TULP1), TULP2 and TULP3. When compared to TULP1 and TULP2, TULP3 has a wider tissue expression and is phylogenetically more similar to Tub than either TULP1 or TULP2. TULP1, expressed specifically in the retina, maps to the chromosomal region known to be involved in retinitis pigmentosa, while TULP2 maps within the minimal interval for the rod-cone dystrophy. TULP3 maps to human chromosome 12p13, and shares 69% homology to mouse TULP3. Human RNA from testis, ovary, thyroid and spinal cord contain highly detectable levels of TULP3 transcripts. In the retina, TULP3 is expressed specifically in the inner nuclear layer and ganglion cell layer. TULP1, TULP2 and TULP3 may comprise a unique family of bipartite transcription factors.

## TULP3 Polyclonal Antibody - Additional Information

**Gene ID** 7289

### Other Names

Tubby-related protein 3, Tubby-like protein 3, TULP3 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=12425](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=12425)), TUBL3

### Target/Specificity

Expressed at high levels in testis, ovaries, thyroid, and spinal chord.

### Dilution

WB~1:1000  
IHC-P~N/A  
IHC-F~N/A  
IF~1:50~200  
ICC~N/A  
E~N/A

### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## TULP3 Polyclonal Antibody - Protein Information

**Name** TULP3 ([HGNC:12425](#))

**Synonyms** TUBL3

### Function

Negative regulator of the Shh signaling transduction pathway: recruited to primary cilia via association with the IFT complex A (IFT- A) and is required for recruitment of G protein-coupled receptor GPR161 to cilia, a promoter of PKA-dependent basal repression machinery in Shh signaling. Binds to phosphorylated inositide (phosphoinositide) lipids. Both IFT-A- and phosphoinositide-binding properties are required to regulate ciliary G protein-coupled receptor trafficking. During adipogenesis, regulates ciliary trafficking of FFAR4 in preadipocytes.

### Cellular Location

Nucleus. Cell membrane. Cell projection, cilium. Cytoplasm. Secreted. Note=Does not have a cleavable signal peptide and is secreted by a non-conventional pathway (By similarity). Translocates from the plasma membrane to the nucleus upon activation of guanine nucleotide-binding protein G(q) subunit alpha

### Tissue Location


Expressed at high levels in testis, ovaries, thyroid, and spinal cord.

## TULP3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TULP3 Polyclonal Antibody - Images



Sample: 293T Cell (human) Lysate at 40 ug  
Primary: Anti-TULP3(bs-11605R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 50 kD  
Observed band size: 50 kD