

PPP1R10 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54589**Specification**

PPP1R10 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O96QC0
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	99058

PPP1R10 Polyclonal Antibody - Additional Information**Gene ID** 5514**Other Names**

Serine/threonine-protein phosphatase 1 regulatory subunit 10, MHC class I region proline-rich protein CAT53, PP1-binding protein of 114 kDa, Phosphatase 1 nuclear targeting subunit, Protein FB19, p99, PPP1R10, CAT53, FB19, PNUTS

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PPP1R10 Polyclonal Antibody - Protein Information**Name** PPP1R10**Synonyms** CAT53, FB19, PNUTS**Function**

Scaffold protein which mediates the formation of the PTW/PP1 phosphatase complex by providing a binding platform to each component of the complex. The PTW/PP1 phosphatase complex plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. Mediates interaction of WDR82 and PPP1CA. Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00649, ECO:0000269|PubMed:9450550}.

Note=Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity)


Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.

PPP1R10 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPP1R10 Polyclonal Antibody - Images

Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;


Incubation: Anti-ppp1R10 Polyclonal Antibody, Unconjugated(bs-11666R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-ppp1R10 Polyclonal Antibody, Unconjugated(bs-11666R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample:

Spleen (Mouse) Lysate at 40 ug

Primary: Anti-PPP1R10 (Bs- 11666R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 99 kD

Observed band size: 99 kD