

UBXD2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54592**Specification**

UBXD2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O92575
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56778

UBXD2 Polyclonal Antibody - Additional Information**Gene ID** 23190**Other Names**

UBX domain-containing protein 4, Erasin, UBX domain-containing protein 2, UBXN4, KIAA0242, UBXD2, UBXDC1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

UBXD2 Polyclonal Antibody - Protein Information**Name** UBXN4**Synonyms** KIAA0242, UBXD2, UBXDC1**Function**

Involved in endoplasmic reticulum-associated protein degradation (ERAD). Acts as a platform to recruit both UBQLN1 and VCP to the ER during ERAD (PubMed:19822669).

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Nucleus envelope. Note=Both the N- and the C-terminus face the cytosol. Also found in the nucleus envelope contiguous to the ER

Tissue Location


Expressed in many tissues, including heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas Accumulates in Alzheimer disease-afflicted brains (at protein level)

UBXD2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UBXD2 Polyclonal Antibody - Images



Sample:

Heart (Mouse) Lysate at 40 ug


Cerebrum (Rat) Lysate at 40 ug

Primary: Anti-UBXD2 (bs-11675R) at 1/1000 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 57 kD

Observed band size: 57 kD



Paraformaldehyde-fixed, paraffin embedded (Human kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UBXD2) Polyclonal Antibody, Unconjugated (bs-11675R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (UBXD2) Polyclonal Antibody, Unconjugated (bs-11675R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.