

ATBF1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54629

Specification

ATBF1 Polyclonal Antibody - Product Information

Application **Primary Accession**

Reactivity Host Clonality Calculated MW **Physical State** Immunogen

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

IHC-P, IHC-F, IF, ICC, E

015911

Rat, Dog, Bovine

Rabbit **Polyclonal** 404 KDa Liquid

KLH conjugated synthetic peptide derived

from human ATBF1

301-400/3703

laG

Buffer

SUBCELLULAR LOCATION

SIMILARITY

SUBUNIT

Post-translational modifications

Important Note

0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Contains 22 C2H2-type zinc fingers. Contains 4 homeobox DNA-binding

domains.

Interacts with FNBP3 (By similarity).

Interacts with PIAS3.

Phosphorylated upon DNA damage,

probably by ATM or ATR.

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

AT-motif binding factor 1 (ATBF1) binds to the AT-rich core sequence element in the human a-fetoprotein enhancer (1). Alternative splicing generates the ATBF1-A and ATBF1-B (2,3). While ATBF1-A contains a 920-amino acid extension at the N-terminus, both ATBF1-A and ATBF1-B contain 4 DNA-binding homeobox domains (2,3). Additionally, ATBF1-A contains 23 zinc finger motifs while ATBF1-B contains 18 zinc finger motifs (1-3). The N-terminal extension unique to ATBF1-A has transcriptional repressor activity (4). In the small intestine, ATBF1-A inhibits expression of the brushborder enzyme aminopeptidase-N through direct binding to the AT motif element (5). Besides functioning in transcription regulation, ATBF1 also functions in ATPase activity (6). ATPase activity associated with ATBF1-A is DNA/RNA-dependent and requires both homeobox domains and zinc finger motifs (6). ATBF1 is highly expressed in spleen and brain tissues (7). The gene encoding human ATBF1 maps to chromosome 16q22.3-q23.1 (8).

ATBF1 Polyclonal Antibody - Additional Information

Gene ID 463



Other Names

Zinc finger homeobox protein 3, AT motif-binding factor 1, AT-binding transcription factor 1, Alpha-fetoprotein enhancer-binding protein, Zinc finger homeodomain protein 3, ZFH-3, ZFHX3, ATBF1, C16orf47 {ECO:0000312|HGNC:HGNC:777}

Target/Specificity

Not found in normal gastric mucosa but found in gastric carcinoma cells (at protein level).

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> ICC~~N/A<br \> E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ATBF1 Polyclonal Antibody - Protein Information

Name ZFHX3

Synonyms ATBF1, C16orf47 {ECO:0000312|HGNC:HGNC:7

Function

Transcriptional regulator which can act as an activator or a repressor. Inhibits the enhancer element of the AFP gene by binding to its AT-rich core sequence. In concert with SMAD-dependent TGF-beta signaling can repress the transcription of AFP via its interaction with SMAD2/3 (PubMed:25105025). Regulates the circadian locomotor rhythms via transcriptional activation of neuropeptidergic genes which are essential for intercellular synchrony and rhythm amplitude in the suprachiasmatic nucleus (SCN) of the brain (By similarity). Regulator of myoblasts differentiation through the binding to the AT-rich sequence of MYF6 promoter and promoter repression (PubMed: 11312261). Down-regulates the MUC5AC promoter in gastric cancer (PubMed: 17330845). In association with RUNX3, up-regulates CDKN1A promoter activity following TGF-beta stimulation (PubMed:20599712). Inhibits estrogen receptor (ESR1) function by selectively competing with coactivator NCOA3 for binding to ESR1 in ESR1-positive breast cancer cells (PubMed:20720010).

Cellular Location

Nucleus. Cytoplasm Note=Translocates from the cytoplasm to the nucleus following TGF-beta stimulation. Expressed in nuclear body (NB)-like dots in the nucleus some of which overlap or closely associate with PML body

Tissue Location

Not found in normal gastric mucosa but found in gastric carcinoma cells (at protein level). Expression is higher in ER- positive breast tumors than ER-negative breast tumors (at protein level).

ATBF1 Polyclonal Antibody - Protocols






Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ATBF1 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ATBF1) Polyclonal Antibody, Unconjugated (bs-11805R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.