

CSDC2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54637

Specification

CSDC2 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>O9Y534</u>

Reactivity Rat, Pig, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 16786

CSDC2 Polyclonal Antibody - Additional Information

Gene ID 27254

Other Names

Cold shock domain-containing protein C2, RNA-binding protein PIPPin, CSDC2, PIPPIN

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CSDC2 Polyclonal Antibody - Protein Information

Name CSDC2

Synonyms PIPPIN

Function


RNA-binding factor which binds specifically to the very 3'- UTR ends of both histone H1 and H3.3 mRNAs, encompassing the polyadenylation signal. Might play a central role in the negative regulation of histone variant synthesis in the developing brain (By similarity).

Cellular Location

Nucleus. Cytoplasm. Note=PIPPin-RNA complexes are located to the nucleus.

CSDC2 Polyclonal Antibody - Protocols







Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CSDC2 Polyclonal Antibody - Images