

ITM2C Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54649

Specification

ITM2C Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NOX7
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ITM2C
Epitope Specificity	3-88/267
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Lysosome membrane.
SIMILARITY	Belongs to the ITM2 family. Contains 1 BRICHOS domain.
SUBUNIT	Interacts with BACE1. Interacts with APP. Interacts with STMN2.
Post-translational modifications	Type I membrane-bound, as well as soluble, furin has a pre-eminent role in ITM2C proteolytic processing. PCSK7 and PCSK5 may also be involved although to a lesser extent. The soluble form of PCSK7 is incapable of processing ITM2C.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Negative regulator of beta amyloid peptide production. May inhibit the processing of APP by blocking its access to alpha-and beta-secretase. Binding to the beta-secretase-cleaved APP C-terminal fragment is negligible, suggesting that ITM2C is a poor gamma-secretase cleavage inhibitor. May play a role in TNF-induced cell death and neuronal differentiation.

ITM2C Polyclonal Antibody - Additional Information

Gene ID 81618

Other Names

Integral membrane protein 2C, Cerebral protein 14, Transmembrane protein BRI3, CT-BRI3, ITM2C, BRI3

Target/Specificity

High levels in the brain, specifically in the cerebral cortex, medulla, amygdala, hippocampus, thalamus, caudate nucleus, cerebellum, olfactory lobe and spinal cord. Very low levels in other organs.

Dilution

```
<span class ="dilution_WB">WB~~1:1000</span><br ><span class = "dilution_IHC-P">IHC-P~~N/A</span><br ><span class = "dilution_IHC-F">IHC-F~~N/A</span><br ><span class = "dilution_IF">IF~~1:50~200</span><br ><span class ="dilution_ICC">ICC~~N/A</span><br ><span class ="dilution_E">E~~N/A</span>
```

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ITM2C Polyclonal Antibody - Protein Information

Name ITM2C

Synonyms BRI3

Function

Negative regulator of amyloid-beta peptide production. May inhibit the processing of APP by blocking its access to alpha- and beta-secretase. Binding to the beta-secretase-cleaved APP C-terminal fragment is negligible, suggesting that ITM2C is a poor gamma-secretase cleavage inhibitor. May play a role in TNF-induced cell death and neuronal differentiation (By similarity).

Cellular Location

Lysosome membrane; Single-pass type II membrane protein. Cell membrane; Single-pass type II membrane protein

Tissue Location

High levels in the brain, specifically in the cerebral cortex, medulla, amygdala, hippocampus, thalamus, caudate nucleus, cerebellum, olfactory lobe and spinal cord. Very low levels in other organs.

ITM2C Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ITM2C Polyclonal Antibody - Images