

**KIRREL3 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54653**

**Specification**

**KIRREL3 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q8IZU9</a>
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	83 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KIRREL3
Epitope Specificity	351-450/778
Isotype	IgG
<b>Purity</b>	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Single-pass type I membrane protein
SIMILARITY	Belongs to the immunoglobulin superfamily. Contains 5 Ig-like C2-type (immunoglobulin-like) domains. Interacts with the C-terminus of NPHS2/podocin. Interacts with CASK. Phosphorylation probably regulates the interaction with NSH2. Phosphorylated at Tyr-605 and Tyr-606 by FYN, leading to GRB2 binding (By similarity). N-glycosylated (By similarity).
SUBUNIT	Note=A chromosomal aberration involving KIRREL3 and CDH15 is found in a patient with severe mental retardation and dysmorphic facial features. Translocation t(11;16)(q24.2;q24). Defects in KIRREL3 are the cause of mental retardation autosomal dominant type 4 (MRD4) [MIM:612581]. Mental retardation is characterized by significantly sub-average general intellectual functioning associated with impairments in adaptive behavior and manifested during the developmental period.
Post-translational modifications	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
DISEASE	
Important Note	

**Background Descriptions**

NEPH2 is a 778 amino acid single-pass type I membrane protein that belongs to the nephrin-like protein family and immunoglobulin superfamily. Expressed in both fetal and adult brain, as well as podocytes of kidney glomeruli, NEPH2 contains five Ig-like C2-type (immunoglobulin-like) domains and is thought to play a role in the hematopoietic supportive capacity of stroma cells. NEPH2 undergoes alternative splicing to produce two isoforms and contains a C-terminal cytoplasmic domain which it uses to interact with Podocin, a podocyte protein involved in ultrafiltration. Defects in the gene encoding NEPH2 are associated with mental retardation autosomal dominant type 4 (MRD4).

## KIRREL3 Polyclonal Antibody - Additional Information

**Gene ID** 84623

### Other Names

Kin of IRRE-like protein 3, Kin of irregular chiasm-like protein 3, Nephrin-like protein 2, Processed kin of IRRE-like protein 3, KIRREL3 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=23204" target="\_blank">HGNC:23204</a>)

### Target/Specificity

Expressed in fetal and adult brain. Also expressed in kidney, specifically in podocytes of kidney glomeruli.

### Dilution

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br /><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br /><span class ="dilution\_IF">IF~~1:50~200</span><br /><span class ="dilution\_ICC">ICC~~N/A</span><br /><span class ="dilution\_E">E~~N/A</span>

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## KIRREL3 Polyclonal Antibody - Protein Information

**Name** KIRREL3 ([HGNC:23204](#))

### Function

Synaptic adhesion molecule required for the formation of target-specific synapses. Required for formation of target-specific synapses at hippocampal mossy fiber synapses. Required for formation of mossy fiber filopodia, the synaptic structures connecting dentate granule and GABA neurons. Probably acts as a homophilic adhesion molecule that promotes trans-cellular interactions and stabilize mossy fiber filopodia contact and subsequent synapse formation. Required for the coalescence of vomeronasal sensory neuron axons. May be involved in the hematopoietic supportive capacity of stroma cells; the secreted extracellular domain is directly responsible for supporting hematopoietic stem cells.

### Cellular Location

Cell membrane; Single-pass type I membrane protein

### Tissue Location

Expressed in fetal and adult brain (PubMed:19012874). Also expressed in kidney, specifically in podocytes of kidney glomeruli (PubMed:12424224). Also expressed in skeletal muscle


(PubMed:25488023).

## KIRREL3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## KIRREL3 Polyclonal Antibody - Images



Blank control(blue):Mouse nephrocytes (fixed with 2% paraformaldehyde (10 min)).

Primary Antibody:Rabbit Anti- KIRREL3 antibody(bs-11864R), Dilution: 1  $\mu$ g in 100  $\mu$ L 1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions );

Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.