

EBF1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54730**Specification****EBF1 Polyclonal Antibody - Product Information**

| | |
|----------------------|--|
| Application | IHC-P, IHC-F, IF, ICC, E |
| Primary Accession | Q9UH73 |
| Reactivity | Rat, Dog, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 64 KDa |
| Physical State | Liquid |
| Immunogen | KLH conjugated synthetic peptide derived from human EBF1 |
| Epitope Specificity | 75-170/591 |
| Isotype | IgG |
| Purity | affinity purified by Protein A |
| Buffer | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| SUBCELLULAR LOCATION | Nucleus (Potential). |
| SIMILARITY | Belongs to the COE family. Contains 1 IPT/TIG domain. |
| SUBUNIT | Forms either a homodimer or a heterodimer with a related family member (By similarity). Interacts with ZNF423 and ZNF521, leading to prevent EBF1 to bind DNA and activate target genes. |
| Important Note | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |

Background Descriptions

Transcriptional activator which recognizes variations of the palindromic sequence 5'-ATTCCCNNGGAATT-3'.

EBF1 Polyclonal Antibody - Additional Information**Gene ID 1879****Other Names**

Transcription factor COE1, O/E-1, OE-1, Early B-cell factor, EBF1, COE1, EBF

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
ICC~~N/A
E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

EBF1 Polyclonal Antibody - Protein Information

Name EBF1

Synonyms COE1, EBF

Function

Key pioneer transcription factor of B-cell specification and commitment (PubMed:27807034). Recognizes variations of the palindromic sequence 5'-ATCCCCNNGGAATT-3'. Operates in a transcription factor network to activate B-cell-specific genes and repress genes associated with alternative cell fates. For instance, positively regulates many B- cell specific genes including BCR or CD40 while repressing genes that direct cells into alternative lineages, including GATA3 and TCF7 for the T-cell lineage. In addition to its role during lymphopoiesis, controls the thermogenic gene program in adipocytes during development and in response to environmental cold (By similarity).

Cellular Location

Nucleus.

EBF1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EBF1 Polyclonal Antibody - Images