

SCN3A Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54739

Specification

SCN3A Polyclonal Antibody - Product Information

IHC-P, IHC-F, IF, ICC, E Application

Primary Accession 09NY46

Reactivity Rat, Pig, Bovine Host **Rabbit** Clonality **Polyclonal** Calculated MW 226 KDa

Physical State Liquid Immunogen KLH conjugated synthetic peptide derived

from human SCN3A

1401-1600/2000 **Epitope Specificity** laG

Isotype **Purity**

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Membrane; Multi-pass membrane protein.

SIMILARITY Belongs to the sodium channel (TC

1.A.1.10) family. Nav1.3/SCN3A subfamily.

Contains 1 IO domain.

SUBUNIT The sodium channel consists of a large

polypeptide and 2-3 smaller ones. This sequence represents a large polypeptide.

Interacts with NEDD4L.

Post-translational modifications May be ubiquitinated by NEDD4L; which

would promote its endocytosis.

Important Note This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

Voltage-gated sodium channels are transmembrane glycoprotein complexes composed of a large alpha subunit with 24 transmembrane domains and one or more regulatory beta subunits. They are responsible for the generation and propagation of action potentials in neurons and muscle. This gene encodes one member of the sodium channel alpha subunit gene family, and is found in a cluster of five alpha subunit genes on chromosome 2. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008].

SCN3A Polyclonal Antibody - Additional Information

Gene ID 6328

Other Names

Sodium channel protein type 3 subunit alpha, Sodium channel protein brain III subunit alpha,



Sodium channel protein type III subunit alpha, Voltage-gated sodium channel subtype III, Voltage-gated sodium channel subunit alpha Nav1.3, SCN3A, KIAA1356, NAC3

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> ICC~~N/A<br \> E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SCN3A Polyclonal Antibody - Protein Information

Name SCN3A (HGNC:10590)

Function

Pore-forming subunit of Nav1.3, a voltage-gated sodium (Nav) channel that directly mediates the depolarizing phase of action potentials in excitable membranes. Navs, also called VGSCs (voltage-gated sodium channels) or VDSCs (voltage-dependent sodium channels), operate by switching between closed and open conformations depending on the voltage difference across the membrane. In the open conformation they allow Na(+) ions to selectively pass through the pore, along their electrochemical gradient. The influx of Na+ ions provokes membrane depolarization, initiating the propagation of electrical signals throughout cells and tissues (PubMed:24157691, PubMed:28235671, PubMed:29466837, PubMed:35277491, PubMed:35277491, In some secretory cell types, it also participates in cell excitability through membrane depolarization and regulates cells responsiveness to stimuli triggering secretion. For instance, it controls the release of serotonin/5-hydroxytryptamine by enterochromaffin cells and is required for both glucagon- and glucose- induced insulin secretion in pancreatic endocrine cells (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Basal cell membrane {ECO:0000250|UniProtKB:A2ASI5}; Multi-pass membrane protein. Note=In enterochromaffin cells, localized highly asymmetrically, almost exclusively at the basal side {ECO:0000250|UniProtKB:A2ASI5}

Tissue Location

Expressed in enterochromaffin cells in both colon and small bowel (at protein level).

SCN3A Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot





• <u>Immunohistochemistry</u>

- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SCN3A Polyclonal Antibody - Images