

KCNE4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54761**Specification**

KCNE4 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	Q8WWG9
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human KCNE4
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
SIMILARITY	Belongs to the potassium channel KCNE family.
SUBUNIT	Associates with KCNQ1/KVLQT1 (Probable).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, isk-related subfamily. This member is a type I membrane protein, and a beta subunit that assembles with a potassium channel alpha-subunit to modulate the gating kinetics and enhance stability of the multimeric complex. This gene is prominently expressed in the embryo and in adult uterus. [provided by RefSeq, Jul 2008].

KCNE4 Polyclonal Antibody - Additional Information**Gene ID** 23704**Other Names**

Potassium voltage-gated channel subfamily E member 4, MinK-related peptide 3, Minimum potassium ion channel-related peptide 3, Potassium channel subunit beta MiRP3, KCNE4

Target/Specificity

Predominantly expressed in embryo and adult uterus. Low expression found in kidney, small intestine, lung and heart.

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
ICC ~ ~ N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KCNE4 Polyclonal Antibody - Protein Information

Name KCNE4 ([HGNC:6244](#))

Function

Ancillary protein that functions as a regulatory subunit of the voltage-gated potassium (Kv) channel complex composed of pore- forming and potassium-conducting alpha subunits and of regulatory beta subunits. KCNE4 beta subunit modulates the gating kinetics and enhances stability of the channel complex (PubMed: [12096056](http://www.uniprot.org/citations/12096056), PubMed: [19687231](http://www.uniprot.org/citations/19687231), PubMed: [20533308](http://www.uniprot.org/citations/20533308), PubMed: [27162025](http://www.uniprot.org/citations/27162025)). Associates with KCNQ1/KVLTQ1 alpha subunit to inhibit potassium currents (PubMed: [12096056](http://www.uniprot.org/citations/12096056), PubMed: [19687231](http://www.uniprot.org/citations/19687231), PubMed: [20533308](http://www.uniprot.org/citations/20533308), PubMed: [27162025](http://www.uniprot.org/citations/27162025)).

Cellular Location

Membrane; Single- pass membrane protein

Tissue Location

Predominantly expressed in embryo and adult uterus. Low expression found in kidney, small intestine, lung and heart

KCNE4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KCNE4 Polyclonal Antibody - Images