

HES2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54843

Specification

HES2 Polyclonal Antibody - Product Information

Application IHC-P Primary Accession O9Y543

Reactivity Rat, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 18470

HES2 Polyclonal Antibody - Additional Information

Gene ID 54626

Other Names

Transcription factor HES-2, Class B basic helix-loop-helix protein 40, bHLHb40, Hairy and enhancer of split 2, HES2, BHLHB40

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HES2 Polyclonal Antibody - Protein Information

Name HES2

Synonyms BHLHB40

Function

Transcriptional repressor of genes that require a bHLH protein for their transcription.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00380, ECO:0000255|PROSITE-ProRule:PRU00981}

Tissue Location

Expressed in placenta, pancreatic cancer, colon cancer with RER, cervical cancer, and in head and neck tumors

HES2 Polyclonal Antibody - Protocols






Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HES2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HES2) Polyclonal Antibody, Unconjugated (bs-12383R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.