

KIF13B Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54845**Specification**

KIF13B Polyclonal Antibody - Product Information

Application	IHC-P, WB, IF
Primary Accession	O9NOT8
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	202789

KIF13B Polyclonal Antibody - Additional Information**Gene ID** 23303**Other Names**

Kinesin-like protein KIF13B, Kinesin-like protein GAKIN, KIF13B, GAKIN, KIAA0639

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

KIF13B Polyclonal Antibody - Protein Information**Name** KIF13B**Synonyms** GAKIN, KIAA0639**Function**

Involved in reorganization of the cortical cytoskeleton. Regulates axon formation by promoting the formation of extra axons. May be functionally important for the intracellular trafficking of MAGUKs and associated protein complexes.

Cellular Location

Cytoplasm, cytoskeleton. Cell projection, axon. Note=accumulates at the distal part of the microtubules in the tips of axons, but not of dendrites

Tissue Location


Ubiquitous.

KIF13B Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KIF13B Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIF13B) Polyclonal Antibody, Unconjugated (bs-12387R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:


Ovary (Rat) Lysate at 40 ug

Primary: Anti-KIF13B (bs-12387R) at 1/500 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 203 kD

Observed band size: 203 kD



Paraformaldehyde-fixed, paraffin embedded (Mouse ovary); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (KIF13B) Polyclonal Antibody, Unconjugated (bs-12387R) at 1:400 overnight at 4°C, followed by a conjugated Goat Anti-Rabbit IgG antibody (bs-0295G-AF488) for 90 minutes, and DAPI for nuclei staining.