

ApoM Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54867**Specification**

ApoM Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O95445
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	21253

ApoM Polyclonal Antibody - Additional Information**Gene ID** 55937**Other Names**

Apolipoprotein M, Apo-M, ApoM, Protein G3a, APOM, G3A, NG20

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ApoM Polyclonal Antibody - Protein Information**Name** APOM**Synonyms** G3A, NG20**Function**

Probably involved in lipid transport. Can bind sphingosine-1- phosphate, myristic acid, palmitic acid and stearic acid, retinol, all- trans-retinoic acid and 9-cis-retinoic acid.

Cellular Location

Secreted. Note=Present in high density lipoprotein (HDL) and to a lesser extent in triglyceride-rich lipoproteins (TGRLP) and low density lipoproteins (LDL)

Tissue Location


Plasma protein. Expressed in liver and kidney.

ApoM Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ApoM Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ApoM) Polyclonal Antibody, Unconjugated (bs-12501R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.