

ARFGAP3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54870**Specification****ARFGAP3 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9NP61
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ARFGAP3
Epitope Specificity	412-516/516
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Golgi apparatus membrane. Also found on peripheral punctate structures likely to be endoplasmic reticulum-Golgi intermediate compartment.
SIMILARITY	Contains 1 Arf-GAP domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

ARFGAP1 (ADP-ribosylation factor GTPase-activating protein 1), ARFGAP2 (ADP-ribosylation factor GTPase-activating protein 2) and ARFGAP3 (ADP-ribosylation factor GTPase-activating protein 3) are GTPase-activating proteins (GAP) that are associated with the Golgi apparatus and interact with ADP-ribosylation factor 1 (ARF). These proteins promote hydrolysis of ARF-bound GTP and are required for the dissociation of coat proteins from Golgi-derived membranes and vesicles. Dissociation of the coat proteins is required for the fusion of these vesicles with target compartments. The activity of ARFGAP1, ARFGAP2 and ARFGAP3 is stimulated by phosphoinositides and inhibited by phosphatidylcholine. The genes encoding ARFGAP1, ARFGAP2 and ARFGAP3 map to human chromosomes 20q13.33, 11p11.2 and 22q13.2, respectively.

ARFGAP3 Polyclonal Antibody - Additional Information**Gene ID** 26286**Other Names**

ADP-ribosylation factor GTPase-activating protein 3, ARF GAP 3, ARFGAP3, ARFGAP1

Target/Specificity

Widely expressed. Highest expression in endocrine glands (pancreas, pituitary gland, salivary gland, and prostate) and testis with a much higher expression in the testis than in the ovary.

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ARFGAP3 Polyclonal Antibody - Protein Information

Name ARFGAP3

Synonyms ARFGAP1

Function

GTPase-activating protein (GAP) for ADP ribosylation factor 1 (ARF1). Hydrolysis of ARF1-bound GTP may lead to dissociation of coatmer from Golgi-derived membranes to allow fusion with target membranes.

Cellular Location

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Note=Also found on peripheral punctate structures likely to be endoplasmic reticulum-Golgi intermediate compartment

Tissue Location

Widely expressed. Highest expression in endocrine glands (pancreas, pituitary gland, salivary gland, and prostate) and testis with a much higher expression in the testis than in the ovary

ARFGAP3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ARFGAP3 Polyclonal Antibody - Images