

## **Argonaute 3 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54874

## **Specification**

## **Argonaute 3 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q9H9G7</u>

Reactivity
Host
Rat, Pig, Dog, Bovine
Rabbit

Clonality Polyclonal Calculated MW 97 KDa Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

laG

from human Argonaute 3/eIF2C3

Epitope Specificity 251-350/860

Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm.

SIMILARITY Belongs to the argonaute family. Ago

subfamily. Contains 1 PAZ domain.

Contains 1 Piwi domain.

SUBUNIT Interacts with EIF4B, IMP8, PRMT5 and

TNRC6B. Interacts with APOBEC3F,

APOBEC3G and APOBEC3H.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

### **Background Descriptions**

This gene encodes a member of the Argonaute family of proteins which play a role in RNA interference. The encoded protein is highly basic, contains a PAZ domain and a PIWI domain, and may play a role in short-interfering-RNA-mediated gene silencing. This gene is located on chromosome 1 in a tandem cluster of closely related family members including argonaute 4 and eukaryotic translation initiation factor 2C, 1. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008]

### **Argonaute 3 Polyclonal Antibody - Additional Information**

### **Gene ID 192669**

### **Other Names**

Protein argonaute-3 {ECO:0000255|HAMAP-Rule:MF\_03032}, Argonaute3 {ECO:0000255|HAMAP-Rule:MF\_03032}, hAgo3, 3.1.26.n2, Argonaute RISC catalytic component 3, Eukaryotic translation initiation factor 2C 3 {ECO:0000255|HAMAP-Rule:MF\_03032}, eIF-2C 3 {ECO:0000255|HAMAP-Rule:MF\_03032}, AGO3,



### EIF2C3

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class = "dilution\_ICC">ICC~~N/A</span><br \> <span class = "dilution\_E">E~~N/A</span>

### **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

## **Argonaute 3 Polyclonal Antibody - Protein Information**

Name AGO3

Synonyms EIF2C3

### **Function**

Required for RNA-mediated gene silencing (RNAi). Binds to short RNAs such as microRNAs (miRNAs) and represses the translation of mRNAs which are complementary to them. Proposed to be involved in stabilization of small RNA derivates (siRNA) derived from processed RNA polymerase III-transcribed Alu repeats containing a DR2 retinoic acid response element (RARE) in stem cells and in the subsequent siRNA- dependent degradation of a subset of RNA polymerase II-transcribed coding mRNAs by recruiting a mRNA decapping complex involving EDC4. Possesses RNA slicer activity but only on select RNAs bearing 5'- and 3'-flanking sequences to the region of guide-target complementarity (PubMed:<a href="http://www.uniprot.org/citations/29040713" target="blank">29040713</a>/a>).

## **Cellular Location**

Cytoplasm, P-body {ECO:0000255|HAMAP- Rule:MF 03032, ECO:0000269|PubMed:16081698}

## **Argonaute 3 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **Argonaute 3 Polyclonal Antibody - Images**