

Argonaute 4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54875**Specification**

Argonaute 4 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9HCK5
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	97 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Argonaute 4/eIF2C4
Epitope Specificity	401-500/861
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
SIMILARITY	Belongs to the argonaute family. Ago subfamily. Contains 1 PAZ domain. Contains 1 Piwi domain. Cellular localization.
SUBUNIT	Interacts with EIF4B, IMP8, PRMT5, TNRC6A and TNRC6B.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

This gene encodes a member of the Argonaute family of proteins which play a role in RNA interference. The encoded protein is highly basic containing PAZ and PIWI domains, and it may play a role in short-interfering-RNA-mediated gene silencing. This gene is located on chromosome 1 in a cluster of closely related family members including argonaute 3, and eukaryotic translation initiation factor 2C, 1. [provided by RefSeq, Jul 2008]

Argonaute 4 Polyclonal Antibody - Additional Information**Gene ID** 192670**Other Names**

Protein argonaute-4 {ECO:0000255|HAMAP-Rule:MF_03033}, Argonaute4 {ECO:0000255|HAMAP-Rule:MF_03033}, hAgo4, Argonaute RISC catalytic component 4, Eukaryotic translation initiation factor 2C 4 {ECO:0000255|HAMAP-Rule:MF_03033}, eIF-2C 4 {ECO:0000255|HAMAP-Rule:MF_03033}, eIF2C 4 {ECO:0000255|HAMAP-Rule:MF_03033}, AGO4, EIF2C4, KIAA1567

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Argonaute 4 Polyclonal Antibody - Protein Information

Name AGO4

Synonyms EIF2C4, KIAA1567

Function

Required for RNA-mediated gene silencing (RNAi). Binds to short RNAs such as microRNAs (miRNAs) and represses the translation of mRNAs which are complementary to them. Lacks endonuclease activity and does not appear to cleave target mRNAs. Also required for RNA-directed transcription and replication of the human hepatitis delta virus (HDV).

Cellular Location

Cytoplasm, P-body {ECO:0000255|HAMAP- Rule:MF_03033, ECO:0000269|PubMed:16081698, ECO:0000269|PubMed:19167051}

Argonaute 4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Argonaute 4 Polyclonal Antibody - Images