

**PFDN5 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54910**

**Specification**

---

**PFDN5 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	<a href="#">Q99471</a>
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17328

**PFDN5 Polyclonal Antibody - Additional Information**

**Gene ID** 5204

**Other Names**

Prefoldin subunit 5, Myc modulator 1, c-Myc-binding protein Mm-1, PFDN5, MM1, PFD5

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**PFDN5 Polyclonal Antibody - Protein Information**

**Name** PFDN5

**Synonyms** MM1, PFD5

**Function**

Binds specifically to cytosolic chaperonin (c-CPN) and transfers target proteins to it. Binds to nascent polypeptide chain and promotes folding in an environment in which there are many competing pathways for nonnative proteins. Represses the transcriptional activity of MYC.

**Cellular Location**

[Isoform 1]: Nucleus. [Isoform 3]: Nucleus.

**Tissue Location**

Highly expressed in pancreas and skeletal muscle and moderately in other tissues

**PFDN5 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**PFDN5 Polyclonal Antibody - Images**