

**VPS11 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54957****Specification**

---

**VPS11 Polyclonal Antibody - Product Information**

|                   |                          |
|-------------------|--------------------------|
| Application       | IHC-P, IHC-F, IF, ICC, E |
| Primary Accession | <a href="#">Q9H270</a>   |
| Reactivity        | Rat, Pig, Dog, Bovine    |
| Host              | Rabbit                   |
| Clonality         | Polyclonal               |
| Calculated MW     | 107837                   |

**VPS11 Polyclonal Antibody - Additional Information****Gene ID** 55823**Other Names**

Vacuolar protein sorting-associated protein 11 homolog, hVPS11, RING finger protein 108, VPS11, RNF108

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
ICC ~ ~ N/A  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**VPS11 Polyclonal Antibody - Protein Information****Name** VPS11**Synonyms** RNF108**Function**

Plays a role in vesicle-mediated protein trafficking to lysosomal compartments including the endocytic membrane transport and autophagic pathways. Believed to act as a core component of the putative HOPS and CORVET endosomal tethering complexes which are proposed to be involved in the Rab5-to-Rab7 endosome conversion probably implicating MON1A/B, and via binding SNAREs and SNARE complexes to mediate tethering and docking events during SNARE-mediated membrane fusion. The HOPS complex is proposed to be recruited to Rab7 on the late endosomal membrane and to regulate late endocytic, phagocytic and autophagic traffic

towards lysosomes. The CORVET complex is proposed to function as a Rab5 effector to mediate early endosome fusion probably in specific endosome subpopulations (PubMed:<a href="http://www.uniprot.org/citations/11382755" target="\_blank">11382755</a>, PubMed:<a href="http://www.uniprot.org/citations/23351085" target="\_blank">23351085</a>, PubMed:<a href="http://www.uniprot.org/citations/24554770" target="\_blank">24554770</a>, PubMed:<a href="http://www.uniprot.org/citations/25266290" target="\_blank">25266290</a>, PubMed:<a href="http://www.uniprot.org/citations/25783203" target="\_blank">25783203</a>). Required for fusion of endosomes and autophagosomes with lysosomes (PubMed:<a href="http://www.uniprot.org/citations/25783203" target="\_blank">25783203</a>). Involved in cargo transport from early to late endosomes and required for the transition from early to late endosomes (PubMed:<a href="http://www.uniprot.org/citations/21148287" target="\_blank">21148287</a>). Involved in the retrograde Shiga toxin transport (PubMed:<a href="http://www.uniprot.org/citations/23593995" target="\_blank">23593995</a>).

#### **Cellular Location**

Endosome. Late endosome membrane; Peripheral membrane protein; Cytoplasmic side. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome {ECO:0000269|PubMed:21148287, ECO:0000305}. Cytoplasmic vesicle. Cytoplasmic vesicle, autophagosome. Cytoplasmic vesicle, clathrin-coated vesicle

#### **Tissue Location**

Ubiquitous. Expression was highest in heart and low in lung

### **VPS11 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **VPS11 Polyclonal Antibody - Images**