

## **VPS26 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54959

# **Specification**

## **VPS26 Polyclonal Antibody - Product Information**

Application WB, IHC-P, IHC-F, IF, ICC

Primary Accession <u>075436</u>

Reactivity
Host
Clonality
Calculated MW
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
38 KDa

Calculated MW 38 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human VPS26

Epitope Specificity 21-120/327 Isotype IaG

Isotype
Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

SUBCELLULAR LOCATION Proclin300 and 50% Glycerol.

Cytoplasm. Endosome membrane.

SIMILARITY Belongs to the VPS26 family.

Important Note

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

# **Background Descriptions**

This gene belongs to a group of vacuolar protein sorting (VPS) genes. The encoded protein is a component of a large multimeric complex, termed the retromer complex, involved in retrograde transport of proteins from endosomes to the trans-Golgi network. The close structural similarity between the yeast and human proteins that make up this complex suggests a similarity in function. Expression studies in yeast and mammalian cells indicate that this protein interacts directly with VPS35, which serves as the core of the retromer complex. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

### **VPS26 Polyclonal Antibody - Additional Information**

#### **Gene ID 9559**

#### **Other Names**

Vacuolar protein sorting-associated protein 26A, Vesicle protein sorting 26A, hVPS26, VPS26A {ECO:0000303|PubMed:30213940, ECO:0000312|HGNC:HGNC:12711}

#### **Dilution**

<span class ="dilution WB">WB~~1:1000/>span class

="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class

="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class

="dilution\_IF">IF $\sim$ 1:50 $\sim$ 200</span><br \><span class ="dilution\_ICC">ICC $\sim$ N/A</span>



### **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **VPS26 Polyclonal Antibody - Protein Information**

Name VPS26A {ECO:0000303|PubMed:30213940, ECO:0000312|HGNC:HGNC:12711}

#### **Function**

Acts as a component of the retromer cargo-selective complex (CSC). The CSC is believed to be the core functional component of retromer or respective retromer complex variants acting to prevent missorting of selected transmembrane cargo proteins into the lysosomal degradation pathway. The recruitment of the CSC to the endosomal membrane involves RAB7A and SNX3. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans- Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX3-retromer mediates the retrograde endosome-to-TGN transport of WLS distinct from the SNX-BAR retromer pathway. The SNX27-retromer is believed to be involved in endosome-to-plasma membrane trafficking and recycling of a broad spectrum of cargo proteins (Probable). The CSC seems to act as recruitment hub for other proteins, such as the WASH complex and TBC1D5 (Probable). Required for retrograde transport of lysosomal enzyme receptor IGF2R (PubMed:<a href="http://www.uniprot.org/citations/15078902">http://www.uniprot.org/citations/15078902</a>"

target="\_blank">15078902</a>, PubMed:<a href="http://www.uniprot.org/citations/15078903" target="\_blank">15078903</a>). Required to regulate transcytosis of the polymeric immunoglobulin receptor (plgR-plgA) (PubMed:<a

href="http://www.uniprot.org/citations/15247922" target="\_blank">15247922</a>). Required for the endosomal localization of WASHC2A (indicative for the WASH complex) (PubMed:<a href="http://www.uniprot.org/citations/22070227" target="\_blank">22070227</a>). Required for the endosomal localization of TBC1D5 (PubMed:<a

href="http://www.uniprot.org/citations/20923837" target="\_blank">20923837</a>). Mediates retromer cargo recognition of SORL1 and is involved in trafficking of SORL1 implicated in sorting and processing of APP (PubMed:<a href="http://www.uniprot.org/citations/22279231" target="\_blank">22279231</a>). Involved in retromer-independent lysosomal sorting of F2R (PubMed:<a href="http://www.uniprot.org/citations/16407403" target="\_blank">16407403</a>). Involved in recycling of ADRB2 (PubMed:<a href="http://www.uniprot.org/citations/21602791" target="\_blank">21602791</a>). Enhances the affinity of SNX27 for PDZ-binding motifs in cargo proteins (By similarity).

### **Cellular Location**

Cytoplasm. Endosome membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:P40336}. Early endosome Note=Localizes to tubular profiles adjacent to endosomes (PubMed:15078903). Predominantly found in early not late endosomes (By similarity). {ECO:0000250|UniProtKB:P40336}

### **VPS26 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation



- Flow Cytomety
- Cell Culture

# **VPS26 Polyclonal Antibody - Images**



# Sample:

A431(Human) Cell Lysate at 40 ug MCF-7(Human) Cell Lysate at 40 ug

Primary: Anti-VPS26 (bs-12768R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 38 kD Observed band size: 38 kD