

### **ZNF202 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54976

# **Specification**

# **ZNF202 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC

Primary Accession <u>095125</u>

Reactivity Rat, Pig, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 74720

# **ZNF202 Polyclonal Antibody - Additional Information**

### **Gene ID** 7753

#### **Other Names**

Zinc finger protein 202, Zinc finger protein with KRAB and SCAN domains 10, ZNF202, ZKSCAN10

### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span> <br \> <span class
="dilution\_IF">IF~~1:50~200</span> <br \> <span class = "dilution\_ICC">ICC~~N/A</span>

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **ZNF202 Polyclonal Antibody - Protein Information**

Name ZNF202

Synonyms ZKSCAN10

### **Function**


Transcriptional repressor that binds to elements found predominantly in genes that participate in lipid metabolism. Among its targets are structural components of lipoprotein particles (apolipoproteins AIV, CIII, and E), enzymes involved in lipid processing (lipoprotein lipase, lecithin cholesteryl ester transferase), transporters involved in lipid homeostasis (ABCA1, ABCG1), and several genes involved in processes related to energy metabolism and vascular disease.

### **Cellular Location**

Nucleus.

# **Tissue Location**

Highly expressed in testis. Also expressed in breast carcinoma cell lines





**ZNF202 Polyclonal Antibody - Protocols** 

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**ZNF202 Polyclonal Antibody - Images**