

ZNF496 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54988**Specification**

ZNF496 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q96IT1
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66908

ZNF496 Polyclonal Antibody - Additional Information**Gene ID** 84838**Other Names**

Zinc finger protein 496, Zinc finger protein with KRAB and SCAN domains 17, ZNF496, ZKSCAN17

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZNF496 Polyclonal Antibody - Protein Information**Name** ZNF496**Synonyms** ZKSCAN17**Function**

DNA-binding transcription factor that can both act as an activator and a repressor.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.


ZNF496 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF496 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF496) Polyclonal Antibody, Unconjugated (bs-12815R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF496) Polyclonal Antibody, Unconjugated (bs-12815R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.