

SYBL1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55002

Specification

SYBL1 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Physical State Immunogen Epitope Specificity Isotype Purity affinity purified by Protein A	WB, IHC-P, IHC-F, IF, ICC <u>P51809</u> Rat, Pig, Bovine Rabbit Polyclonal 25 KDa Liquid KLH conjugated synthetic peptide derived from human SYBL1 1-100/220 IgG
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02%
SUBCELLULAR LOCATION	Proclin300 and 50% Glycerol. Cytoplasmic vesicle > secretory vesicle membrane. Golgi apparatus > trans-Golgi network membrane. Late endosome membrane. Lysosome membrane. Endoplasmic reticulum membrane. Cytoplasmic vesicle > phagosome membrane.
SIMILARITY	Belongs to the synaptobrevin family. Contains 1 longin domain. Contains 1 v-SNARE coiled-coil homology domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	
FLI53045, FLI53762, FLI54296, HCNC+11486, OTTHUMP0000024258, OTTHUMP0000024259,	

FLJ53045; FLJ53762; FLJ54296; HGNC:11486; OTTHUMP00000024258; OTTHUMP00000024259; OTTHUMP00000225953; SYBL 1; SYBL1; Synaptobrevin like 1; Synaptobrevin-like protein 1; Tetanus insensitive VAMP; Tetanus neurotoxin insensitive VAMP; Tetanus-insensitive VAMP; TI VAMP; Ti-VAMP; TIVAMP; VAMP-7; VAMP7; VAMP7_HUMAN

SYBL1 Polyclonal Antibody - Additional Information

Gene ID 6845

Other Names Vesicle-associated membrane protein 7, VAMP-7, Synaptobrevin-like protein 1, Tetanus-insensitive VAMP, Ti-VAMP, VAMP7, SYBL1



Target/Specificity Detected in all tissues tested.

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SYBL1 Polyclonal Antibody - Protein Information

Name VAMP7

Synonyms SYBL1

Function

Involved in the targeting and/or fusion of transport vesicles to their target membrane during transport of proteins from the early endosome to the lysosome. Required for heterotypic fusion of late endosomes with lysosomes and homotypic lysosomal fusion. Required for calcium regulated lysosomal exocytosis. Involved in the export of chylomicrons from the endoplasmic reticulum to the cis Golgi. Required for exocytosis of mediators during eosinophil and neutrophil degranulation, and target cell killing by natural killer cells. Required for focal exocytosis of late endocytic vesicles during phagosome formation.

Cellular Location

Cytoplasmic vesicle, secretory vesicle membrane; Single-pass type IV membrane protein Golgi apparatus, trans-Golgi network membrane; Single- pass type IV membrane protein. Late endosome membrane; Single-pass type IV membrane protein Lysosome membrane; Single-pass type IV membrane protein. Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Cytoplasmic vesicle, phagosome membrane; Single-pass type IV membrane protein. Synapse, synaptosome. Note=In immature neurons expression is localized in vesicular structures in axons and dendrites while in mature neurons it is localized to the somatodendritic region Colocalizes with LAMP1 in kidney cells. Localization to the endoplasmic reticulum membrane was observed in the intestine but not in liver or kidney (By similarity).

Tissue Location Detected in all tissues tested.

SYBL1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry



Immunofluorescence

- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

SYBL1 Polyclonal Antibody - Images



Sample:

Liver (Mouse) Lysate at 40 ug Primary: Anti-SYBL1 (bs-12852R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 25 kD Observed band size: 26 kD