

## **TCRGC2 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55013

# **Specification**

# **TCRGC2 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession
Host
Clonality
Calculated MW
Physical State
P03986
Rabbit
Polyclonal
22 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human TRGC2

Epitope Specificity 61-160/189

Isotype IgG
Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Membrane; Single-pass membrane protein

(Potential).

SIMILARITY Contains 1 Ig-like (immunoglobulin-like)

domain.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

#### **Background Descriptions**

affinity purified by Protein A

T cell receptors recognize foreign antigens which have been processed as small peptides and bound to major histocompatibility complex (MHC) molecules at the surface of antigen presenting cells (APC). Each T cell receptor is a dimer consisting of one alpha and one beta chain or one delta and one gamma chain. In a single cell, the T cell receptor loci are rearranged and expressed in the order delta, gamma, beta, and alpha. If both delta and gamma rearrangements produce functional chains, the cell expresses delta and gamma. If not, the cell proceeds to rearrange the beta and alpha loci. This region represents the germline organization of the T cell receptor gamma locus. The gamma locus includes V (variable), J (joining), and C (constant) segments. During T cell development, the gamma chain is synthesized by a recombination event at the DNA level joining a V segment with a | segment; the C segment is later joined by splicing at the RNA level. Recombination of many different V segments with several I segments provides a wide range of antigen recognition. Additional diversity is attained by junctional diversity, resulting from the random addition of nucleotides by terminal deoxynucleotidyltransferase. Several V segments of the gamma locus are known to be incapable of encoding a protein and are considered pseudogenes. Somatic rearrangement of the gamma locus has been observed in T cells derived from patients with T cell leukemia and ataxia telangiectasia. [provided by RefSeq, Jul 2008]

### **TCRGC2 Polyclonal Antibody - Additional Information**

**Other Names** 



T cell receptor gamma constant 2 {ECO:0000303|Ref.4}, TRGC2 {ECO:0000303|Ref.4}

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class = "dilution\_ICC">ICC~~N/A</span><br \> <span class = "dilution\_E">E~~N/A</span>

#### **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# **TCRGC2 Polyclonal Antibody - Protein Information**

Name TRGC2 {ECO:0000303|Ref.4}

#### **Function**

Constant region of T cell receptor (TR) gamma chain that participates in the antigen recognition (PubMed: <a href="http://www.uniprot.org/citations/24600447" target="blank">24600447</a>). Gamma-delta TRs recognize a variety of self and foreign non-peptide antigens frequently expressed at the epithelial boundaries between the host and external environment, including endogenous lipids presented by MH-like protein CD1D and phosphoantigens presented by butyrophilin-like molecule BTN3A1. Upon antigen recognition induces rapid, innate-like immune responses involved in pathogen clearance and tissue repair (PubMed: <a href="http://www.uniprot.org/citations/23348415" target=" blank">23348415</a>, PubMed:<a href="http://www.uniprot.org/citations/28920588" target="blank">28920588</a>). Binding of gamma-delta TR complex to antigen triggers phosphorylation of immunoreceptor tyrosine-based activation motifs (ITAMs) in the CD3 chains by the LCK and FYN kinases, allowing the recruitment, phosphorylation, and activation of ZAP70 that facilitates phosphorylation of the scaffolding proteins LCP2 and LAT. This lead to the formation of a supramolecular signalosome that recruits the phospholipase PLCG1, resulting in calcium mobilization and ERK activation, ultimately leading to T cell expansion and differentiation into effector cells (PubMed: <a href="http://www.uniprot.org/citations/25674089" target=" blank">25674089</a>). Gamma-delta TRs are produced through somatic rearrangement of a limited repertoire of variable (V), diversity (D), and joining (J) genes. The potential diversity of gamma-delta TRs is conferred by the unique ability to rearrange (D) genes in tandem and to utilize all three reading frames. The combinatorial diversity is considerably increased by the sequence exonuclease trimming and random nucleotide (N) region additions which occur during the V-(D)-J rearrangements (PubMed:<a href="http://www.uniprot.org/citations/24387714" target=" blank">24387714</a>).

## **Cellular Location**

Cell membrane.

### **TCRGC2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation





- Flow CytometyCell Culture

**TCRGC2 Polyclonal Antibody - Images**