



CTNS Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55022

Specification

CTNS Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

Epitope Specificity Isotype **Purity** affinity purified by Protein A

Buffer

SUBCELLULAR LOCATION

SIMILARITY

DISEASE

WB, IHC-P, IHC-F, IF, ICC, E

060931

Rat

Rabbit

Polyclonal

42 KDa

Liquid

KLH conjugated synthetic peptide derived from human CTNS/Cystinosin

231-330/367

laG

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. Lysosome membrane; Multi-pass membrane protein.

Belongs to the cystinosin family. Contains 2 PQ-loop domains.

Defects in CTNS are the cause of cystinosis nephropathic type (CTNS) [MIM:219800]. It is a form of cystinosis, a lysosomal storage disease due to defective transport of cystine across the lysosomal membrane. This results in cystine accumulation and crystallization in the cells causing widespread tissue damage. The classical nephropathic form has onset in the first vear of life and is characterized by a polyuro-polydipsic syndrome, marked height-weight growth delay, generalized impaired proximal tubular reabsorptive capacity, with severe fluid-electrolyte balance alterations, renal failure, ocular symptoms and other systemic complications. Defects in CTNS are the cause of cystinosis adult non-nephropathic type (CTNSANN) [MIM:219750]. It is a form of cystinosis, a lysosomal storage disease due to defective transport of cystine across the lysosomal membrane. This results in cystine accumulation and crystallization in the cells causing widespread tissue damage. Cystinosis



adult non-nephropathic type is characterized by ocular features and a benigne course. Patients manifest mild photophobia due to conjunctival and corneal cystine crystals. Defects in CTNS are the cause of cystinosis late-onset juvenile or adolescent nephropathic type (CTNSJAN) [MIM:219900]. It is a form of cystinosis, a lysosomal storage disease due to defective transport of cystine across the lysosomal membrane. This results in cystine accumulation and crystallization in the cells causing widespread tissue damage. Late-onset juvenile or adolescent nephropathic cystinosis manifests itself first at age 10 to 12 years with proteinuria due to glomerular damage rather than with the manifestations of tubular damage that occur first in infantile cystinosis. There is no excess amino aciduria and stature is normal. Photophobia, late development of pigmentary retinopathy, and chronic headaches are features. This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Important Note

Background Descriptions

Cystinosis is an autosomal recessive disorder resulting from defective lysosomal transport of cystine and present at birth as a failure to thrive, rickets and proximal renal tubular acidosis. The human CTNS gene on chromosome 17p13 encodes the protein Cystinosin, and mutations in CTNS are responsible for nephropathic cystinosis. The CTNS promoter contains an Sp1 binding element. Cystinosin is an integral membrane protein containing 7 transmembrane domains that functions as a H+-driven transporter responsible for cystine export from lysosomes. In humans, Cystinosin is expressed abundantly in pancreas, kidney (mature and fetal), and skeletal muscle. The mouse homolog to CTNS encodes a protein which is expressed in all tissues except skeletal muscle. In the cell, Cystinosin co-localizes with LAMP-2 to lysosomes. A C-terminal GYDQL sorting motif within Cystinosin is critical for lysosomal localization.

CTNS Polyclonal Antibody - Additional Information

Gene ID 1497

Other Names

Cystinosin, CTNS {ECO:0000303|PubMed:9537412, ECO:0000312|HGNC:HGNC:2518}

Target/Specificity

Strongly expressed in pancreas, kidney (adult and fetal) and in skeletal muscle. Expressed at lower levels in placenta and heart. Weakly expressed in lung, liver and brain (adult and fetal).

Dilution

- WB~~1:1000/>span class
- ="dilution IHC-P">IHC-P~~N/A<br \><span class
- ="dilution IHC-F">IHC-F~~N/A<br \><span class
- ="dilution IF">IF \sim 1:50 \sim 200
span class ="dilution ICC">ICC \sim N/A
br



\>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CTNS Polyclonal Antibody - Protein Information

Name CTNS {ECO:0000303|PubMed:9537412, ECO:0000312|HGNC:HGNC:2518}

Function

Cystine/H(+) symporter that mediates export of cystine, the oxidized dimer of cysteine, from lysosomes (PubMed: 11689434, PubMed:15128704, PubMed:18337546, PubMed:22232659, PubMed:29467429, PubMed:33208952, PubMed:36113465). Plays an important role in melanin synthesis by catalyzing cystine export from melanosomes, possibly by inhibiting pheomelanin synthesis (PubMed:22649030). In addition to cystine export, also acts as a positive regulator of mTORC1 signaling in kidney proximal tubular cells, via interactions with components of the v-ATPase and Ragulator complexes (PubMed:36113465). Also involved in small GTPase-regulated vesicle trafficking and lysosomal localization of LAMP2A, independently of cystine transporter activity (By similarity).

Cellular Location

[Isoform 1]: Lysosome membrane; Multi-pass membrane protein. Melanosome membrane; Multi-pass membrane protein. Note=AP-3 complex is required for localization to the lysosome.

Tissue Location

Strongly expressed in pancreas, kidney (adult and fetal), skeletal muscle, melanocytes and keratinocytes (PubMed:22649030). Expressed at lower levels in placenta and heart Weakly expressed in lung, liver and brain (adult and fetal) (PubMed:22649030).

CTNS Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CTNS Polyclonal Antibody - Images