

**DNAJC9 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55041****Specification****DNAJC9 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q8WXX5</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29910

**DNAJC9 Polyclonal Antibody - Additional Information****Gene ID** 23234**Other Names**

DnaJ homolog subfamily C member 9, HDJC9, DnaJ protein SB73, DNAJC9

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br \><span class = "dilution\_IF">IF~~1:50~200</span><br \><span class = "dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_E">E~~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**DNAJC9 Polyclonal Antibody - Protein Information****Name** DNAJC9**Function**

Acts as a dual histone chaperone and heat shock co-chaperone (PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>). As a histone chaperone, forms a co-chaperone complex with MCM2 and histone H3-H4 heterodimers; and may thereby assist MCM2 in histone H3-H4 heterodimer recognition and facilitate the assembly of histones into nucleosomes (PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>). May also act as a histone co-chaperone together with TONSL (PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>). May recruit histone chaperones ASF1A, NASP and SPT2 to histone H3-H4 heterodimers (PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>).

Also plays a role as co-chaperone of the HSP70 family of molecular chaperone proteins, such as HSPA1A, HSPA1B and HSPA8 (PubMed:<a href="http://www.uniprot.org/citations/17182002" target="\_blank">17182002</a>, PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>). As a co-chaperone, may play a role in the recruitment of HSP70-type molecular chaperone machinery to histone H3-H4 substrates, thereby maintaining the histone structural integrity (PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>). Exhibits activity to assemble histones onto DNA in vitro (PubMed:<a href="http://www.uniprot.org/citations/33857403" target="\_blank">33857403</a>).

#### Cellular Location

Nucleus. Cytoplasm. Cell membrane. Note=Predominantly nuclear. Translocates to the cytoplasm and membrane after heat shock

#### Tissue Location


Expressed in heart, placenta, liver, skeletal muscle, kidney, pancreas, thymus, ovary, colon and peripheral blood

### DNAJC9 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### DNAJC9 Polyclonal Antibody - Images



#### Sample:


Thymus (Mouse) Lysate at 40 ug

Primary: Anti- DNAJC9 (bs-13024R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 30 kD

Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNAJC9) Polyclonal Antibody, Unconjugated (bs-13024R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.