

# ENPP6 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55056

## Specification

# **ENPP6 Polyclonal Antibody - Product Information**

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC-P, IHC-F, IF, ICC, E <u>O6UWR7</u> Rat, Pig, Dog, Bovine Rabbit Polyclonal 50241

## **ENPP6 Polyclonal Antibody - Additional Information**

Gene ID 133121

**Other Names** 

Glycerophosphocholine cholinephosphodiesterase ENPP6, GPC-Cpde, 3.1.4.-, 3.1.4.38, Choline-specific glycerophosphodiester phosphodiesterase, Ectonucleotide pyrophosphatase/phosphodiesterase family member 6, E-NPP 6, NPP-6, ENPP6 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=23409" target="\_blank">HGNC:23409</a>)

Dilution

<span class ="dilution\_WB">WB~~1:1000</span><br \><span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class ="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class ="dilution E">E~~N/A</span>

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **ENPP6 Polyclonal Antibody - Protein Information**

Name ENPP6 (HGNC:23409)

### Function

Choline-specific glycerophosphodiesterase that hydrolyzes glycerophosphocholine (GPC) and lysophosphatidylcholine (LPC) and contributes to supplying choline to the cells (PubMed:<a href="http://www.uniprot.org/citations/15788404" target="\_blank">15788404</a>). Has a preference for LPC with short (12:0 and 14:0) or polyunsaturated (18:2 and 20:4) fatty acids. In vitro, hydrolyzes only choline-containing lysophospholipids, such as sphingosylphosphorylcholine



(SPC), platelet- activating factor (PAF) and lysoPAF, but not other lysophospholipids (By similarity).

**Cellular Location** 

Cell membrane; Lipid-anchor, GPI-anchor. Note=A small amount of the protein may be found in the extracellular milieu

**Tissue Location** 

Predominantly expressed in kidney and brain. In the kidney, expressed specifically in the proximal tubules and thin descending limbs of Henle (at protein level)

## **ENPP6 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### **ENPP6 Polyclonal Antibody - Images**



Tissue/cell: Rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-ENPP6 Polyclonal Antibody, Unconjugated(bs-13076R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining





Sample:MCF-7 Cell (Human) Lysate at 40 ug Primary: Anti-ENPP6(bs-13076R)at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 46kD Observed band size: 50kD



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ENPP6) Polyclonal Antibody, Unconjugated (bs-13076R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.