

ERp72 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55060**Specification****ERp72 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	P13667
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	71 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ERp72
Epitope Specificity	451-550/645
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Endoplasmic reticulum lumen. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
SIMILARITY	Belongs to the protein disulfide isomerase family. Contains 3 thioredoxin domains.
SUBUNIT	Part of a large chaperone multiprotein complex comprising DNAJB11, HSP90B1, HSPA5, HYOU, PDIA2, PDIA4, PDIA6, PPIB, SDF2L1, UGT1A1 and very small amounts of ERP29, but not, or at very low levels, CALR nor CANX.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Mammals defend themselves against intracellular pathogens through presentation of cytoplasmically derived short pathogenic peptides to the cell surface of cytotoxic T lymphocytes, which subsequently leads to cytotoxic events with respect to the affected cell. Antigen presentation is mediated by major histocompatibility complex (MHC) class I molecules, which bind and coordinate short pathogenic peptides. The proper folding and assembly of MHC class I molecules in the endoplasmic reticulum (ER) involve a number of components. MHC class I molecules assemble in the ER with chaperones before binding to the transporter associated with antigen processing (TAP) protein. ERp57 is a component of the MHC class I pathway that appears to interact with MHC class I molecules before they associate with TAP. ERp72, also designated protein disulfide-isomerase A4, is involved in the catalysis of protein -S-S- bond rearrangement. ERp57 and ERp72 may act as proteases, protein disulfide isomerases, phospholipases or a combination of these.

ERp72 Polyclonal Antibody - Additional Information

Gene ID 9601

Other Names

Protein disulfide-isomerase A4, 5.3.4.1, Endoplasmic reticulum resident protein 70, ER protein 70, ERp70, Endoplasmic reticulum resident protein 72, ER protein 72, ERp-72, ERp72, PDIA4, ERP70, ERP72

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ERp72 Polyclonal Antibody - Protein Information

Name PDIA4

Synonyms ERP70, ERP72

Cellular Location

Endoplasmic reticulum lumen. Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065)

ERp72 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ERp72 Polyclonal Antibody - Images