

FbxL4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55077**Specification**

FbxL4 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9UKA2
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70097

FbxL4 Polyclonal Antibody - Additional Information**Gene ID** 26235**Other Names**

F-box/LRR-repeat protein 4, F-box and leucine-rich repeat protein 4, F-box protein FBL4/FBL5, FBXL4, FBL4, FBL5

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FbxL4 Polyclonal Antibody - Protein Information**Name** FBXL4**Synonyms** FBL4, FBL5**Cellular Location**

Cytoplasm. Nucleus. Mitochondrion.

Tissue Location

Expressed in heart, kidney, liver, lung, pancreas, and placenta, but not in skeletal muscle


FbxL4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FbxL4 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Human lung carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FbxL4) Polyclonal Antibody, Unconjugated (bs-13166R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FbxL4) Polyclonal Antibody, Unconjugated (bs-13166R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.