

FNIP1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55088

Specification

FNIP1 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>Q8TF40</u>

Reactivity Rat, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 130 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human FNIP1

Epitope Specificity 101-200/1166

Isotype

Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm.

SIMILARITY Belongs to the FNIP family.
SUBUNIT Forms homomultimers and

heteromultimers with FNIP2. Interacts with FLCN (via C-terminus), with HSPCA and with the PRKAA1, PRKAB1 and PRKAG1 subunits of 5'-AMP-activated protein kinase (AMPK). Phosphorylated FLCN and

AMPK are preferentially bound.

Post-translational modifications Phosphorylated by AMPK.

Important Note

This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

May be involved in energy and/or nutrient sensing through the AMPK and mTOR signaling pathways. May regulate phosphorylation of RPS6KB1.

FNIP1 Polyclonal Antibody - Additional Information

Gene ID 96459

Other Names

Folliculin-interacting protein 1, FNIP1 {ECO:0000303|PubMed:17028174,

ECO:0000312|HGNC:HGNC:29418}

Target/Specificity

Strong expression is found in the heart, liver placenta, muscle, nasal mucosa, salivary gland and



uvula and moderate expression in kidney and lung. Higher levels detected in clear cell renal cell carcinoma (RCC) and chromophobe RCC than in normal kidney tissue.

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> ICC~~N/A<br \> E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.


FNIP1 Polyclonal Antibody - Protein Information

Name FNIP1 {ECO:0000303|PubMed:17028174, ECO:0000312|HGNC:HGNC:29418}

Function

Binding partner of the GTPase-activating protein FLCN: involved in the cellular response to amino acid availability by regulating the non-canonical mTORC1 signaling cascade controlling the MiT/TFE factors TFEB and TFE3 (PubMed: 17028174, PubMed:18663353, PubMed:24081491, PubMed:37079666). Required to promote FLCN recruitment to lysosomes and interaction with Rag GTPases, leading to activation of the non-canonical mTORC1 signaling (PubMed:24081491). In low-amino acid conditions, component of the lysosomal folliculin complex (LFC) on the membrane of lysosomes, which inhibits the GTPase-activating activity of FLCN, thereby inactivating mTORC1 and promoting nuclear translocation of TFEB and TFE3 (By similarity). Upon amino acid restimulation, disassembly of the LFC complex liberates the GTPase-activating activity of FLCN, leading to activation of mTORC1 and subsequent inactivation of TFEB and TFE3 (PubMed:37079666). Together with FLCN, regulates autophagy: following phosphorylation by ULK1, interacts with GABARAP and promotes autophagy (PubMed:25126726). In addition to its role in mTORC1 signaling, also acts as a co-chaperone of HSP90AA1/Hsp90: following gradual phosphorylation by CK2, inhibits the ATPase activity of HSP90AA1/Hsp90, leading to activate both kinase and non-kinase client proteins of HSP90AA1/Hsp90 (PubMed: 27353360, PubMed:30699359). Acts as a scaffold to load client protein FLCN onto HSP90AA1/Hsp90 (PubMed: 27353360). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:27353360). Also acts as a core component of the reductive stress response by inhibiting activation of mitochondria in normal conditions: in response to reductive stress, the conserved Cys degron is reduced, leading to recognition and polyubiquitylation by the CRL2(FEM1B) complex, followed by proteasomal (By similarity). Required for B-cell development (PubMed:32905580).

Cellular Location



abcepta

Lysosome membrane. Cytoplasm, cytosol. Note=Localizes to lysosome membrane in amino acid-depleted conditions and relocalizes to the cytosol upon refeeding (PubMed:29848618). Colocalizes with FLCN in the cytoplasm (PubMed:18663353).

Tissue Location


Strong expression is found in the heart, liver placenta, muscle, nasal mucosa, salivary gland and uvula and moderate expression in kidney and lung. Higher levels detected in clear cell renal cell carcinoma (RCC) and chromophobe RCC than in normal kidney tissue. Expressed in peripheral blood mononuclear cells (PubMed:32181500).

FNIP1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FNIP1 Polyclonal Antibody - Images



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min;

Incubation: Anti-FNIP1 Polyclonal Antibody, Unconjugated(bs-13193R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining