

GABPA Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55110

Specification

GABPA Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession Q06546

Reactivity
Host
Clonality
Calculated MW
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
51 KDa

Physical State
Liquid
Immunogen
KLH conjugated synthetic peptide derived

from human GABPA

Epitope Specificity 51-150/454

Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus.

SIMILARITY Belongs to the ETS family. Contains 1 ETS

DNA-binding domain. Contains 1 PNT

(pointed) domain.

SUBUNIT Heterotetramer of two alpha and two beta

subunits.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

The transcription factor GA-binding protein (GABP) is composed of two subunits, the Ets-related GABP-alpha and a GABP-alpha-associated subunit, GABP beta. GABP alpha binds to a specific DNA sequence and GABP beta exists as b1 and b2 splice variants that differ in their C-termini. In primary neuronal cultures, GABP beta is expressed in both the cytoplasm and the nucleus, whereas GABP alpha is expressed mainly in the nucleus. GABP is constitutively expressed as either a GABP alpha beta heterodimer or a GABP alpha b heterotetramer, both of which can modify GABP-dependent transcription in vitro and in vivo. The GABP alpha beta tetrameric complex performs many different functions, such as stimulating transcription of the adenovirus E4 gene, differentially activating BRCA1 expression in human breast cell lines, potentiating Tat-mediated activation of long terminal repeat promoter transcription and viral replication in certain cell types, acting as a coordinator of mitochrondrial and nuclear transcription for cytochrome oxidase in neurons and assisting in the regulation of rpL32 gene transcription.

GABPA Polyclonal Antibody - Additional Information

Gene ID 2551

Other Names



GA-binding protein alpha chain, GABP subunit alpha, Nuclear respiratory factor 2 subunit alpha, Transcription factor E4TF1-60, GABPA, E4TF1A

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

GABPA Polyclonal Antibody - Protein Information

Name GABPA

Synonyms E4TF1A

Function

Transcription factor capable of interacting with purine rich repeats (GA repeats). Positively regulates transcription of transcriptional repressor RHIT/ZNF205 (PubMed:22306510).

Cellular Location

Nucleus.


GABPA Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GABPA Polyclonal Antibody - Images





Paraformaldehyde-fixed, paraffin embedded (Rat kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GABPA) Polyclonal Antibody, Unconjugated (bs-13261R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

293T Cell (Human) Lysate at 30 ug

Primary: Anti- GABPA (bs-13261R)at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 51kD Observed band size: 51kD