

**GALNT11 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55116**

**Specification**

**GALNT11 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">Q8NCW6</a>
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	69 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GALNT11/GaINAc-T11 401-500/608
Epitope Specificity	IgG
Isotype	
<b>Purity</b>	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Golgi apparatus membrane.
SIMILARITY	Belongs to the glycosyltransferase 2 family. GaINAc-T subfamily. Contains 1 ricin B-type lectin domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

The UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase (GaINAc-T) family of enzymes are substrate-specific proteins that catalyze the transfer of GaINAc (N-acetylgalactosamine) to serine and threonine residues onto various proteins, thereby initiating mucin-type O-linked glycosylation in the Golgi apparatus. GaINAc-T11 (Polypeptide N-acetylgalactosaminyltransferase 11), also known as UDP-GaINAc:polypeptide N-acetylgalactosaminyltransferase 11, is a 608 amino acid protein that catalyzes glycosylation of Muc1, Muc4.1 and EA2, though it does not display enzymatic preference for erythropoitein. The N-terminal domain is involved in substrate binding and manganese coordination, while the C-terminal domain is involved in UDP-Gal binding and catalytic reaction. GaINAc-T11 is highly expressed in kidney tubules, though it is not expressed in glomeruli. There are two isoforms of GaINAc-T11 that are produced as a result of alternative splicing events.

**GALNT11 Polyclonal Antibody - Additional Information**

**Gene ID 63917**

**Other Names**

Polypeptide N-acetylgalactosaminyltransferase 11, 2.4.1.41, Polypeptide GaINAc transferase 11, GaINAc-T11, pp-GaNTase 11, Protein-UDP acetylgalactosaminyltransferase 11,

## UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase 11, GALNT11

### Target/Specificity

Highly expressed in kidney. Expressed at intermediate level in brain, heart and skeletal muscle. Weakly expressed other tissues. In kidney, it is strongly expressed in tubules but not expressed in glomeruli.

### Dilution

<span class ="dilution\_WB">WB~~1:1000</span><br /><span class ="dilution\_IHC-P">IHC-P~~N/A</span><br /><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br /><span class ="dilution\_IF">IF~~1:50~200</span><br /><span class ="dilution\_ICC">ICC~~N/A</span><br /><span class ="dilution\_E">E~~N/A</span>

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## GALNT11 Polyclonal Antibody - Protein Information

### Name GALNT11

### Function

Polypeptide N-acetylgalactosaminyltransferase that catalyzes the initiation of protein O-linked glycosylation and is involved in left/right asymmetry by mediating O-glycosylation of NOTCH1. O-glycosylation of NOTCH1 promotes activation of NOTCH1, modulating the balance between motile and immotile (sensory) cilia at the left-right organiser (LRO). Polypeptide N-acetylgalactosaminyltransferases catalyze the transfer of an N-acetyl-D-galactosamine residue to a serine or threonine residue on the protein receptor. Displays the same enzyme activity toward MUC1, MUC4, and EA2 than GALNT1. Not involved in glycosylation of erythropoietin (EPO).

### Cellular Location

Golgi apparatus membrane; Single-pass type II membrane protein

### Tissue Location


Highly expressed in kidney. Expressed at intermediate level in brain, heart and skeletal muscle. Weakly expressed other tissues. In kidney, it is strongly expressed in tubules but not expressed in glomeruli.

## GALNT11 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## GALNT11 Polyclonal Antibody - Images




Sample:Raji(human) Cell Lysate at 40 ug

Primary: Anti-GALNT11(bs-13271R)at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 69kD

Observed band size: 69kD




Sample:siha(human) Cell Lysate at 40 ug

Primary: Anti-GALNT11(bs-13271R)at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 69kD

Observed band size: 69kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GALNT11) Polyclonal Antibody, Unconjugated (bs-13271R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.