

GAPT Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55120**Specification****GAPT Polyclonal Antibody - Product Information**

| | |
|-------------------|------------------------------|
| Application | WB, IHC-P, IHC-F, IF, ICC, E |
| Primary Accession | Q8N292 |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 17883 |

GAPT Polyclonal Antibody - Additional Information**Gene ID** 202309**Other Names**

Protein GAPT, Growth factor receptor-bound protein 2-binding adapter protein, transmembrane, GAPT, C5orf29

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

GAPT Polyclonal Antibody - Protein Information**Name** GAPT**Synonyms** C5orf29**Function**

Negatively regulates B-cell proliferation following stimulation through the B-cell receptor. May play an important role in maintenance of marginal zone (MZ) B-cells (By similarity).

Cellular Location

Cell membrane; Single-pass membrane protein

Tissue Location

Highly expressed in spleen and PBL, detected at lower levels in thymus, and undetectable in all other tissues tested Also expressed in various B-cell lines, monocytic cell line THP-1 and NK-like cell line YT, but not in T-cell line Jurkat or HeLa cells

GAPT Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GAPT Polyclonal Antibody - Images