

GCAP2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55128

Specification

GCAP2 Polyclonal Antibody - Product Information

Application

Primary Accession

Reactivity

Host

Clonality

Calculated MW

WB, IHC-P, IHC-F, IF, ICC

O9UMX6

Rat, Bovine

Rabbit

Polyclonal

23420

GCAP2 Polyclonal Antibody - Additional Information

Gene ID 2979

Other Names

Guanylyl cyclase-activating protein 2, GCAP 2, Guanylate cyclase activator 1B, GUCA1B, GCAP2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

GCAP2 Polyclonal Antibody - Protein Information

Name GUCA1B

Synonyms GCAP2

Function


Stimulates two retinal guanylyl cyclases (GCs) GUCY2D and GUCY2F when free calcium ions concentration is low, and inhibits GUCY2D and GUCY2F when free calcium ions concentration is elevated (By similarity). This Ca(2+)-sensitive regulation of GCs is a key event in recovery of the dark state of rod photoreceptors following light exposure (By similarity). May be involved in cone photoreceptor response and recovery of response in bright light (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P51177}; Lipid-anchor {ECO:0000250|UniProtKB:P51177}. Photoreceptor inner segment. Cell projection, cilium, photoreceptor outer segment Note=Subcellular location is not affected by light or dark conditions {ECO:0000250|UniProtKB:Q8VBV8}

Tissue Location

In the retina, it is expressed in cone and rod photoreceptor cells.





GCAP2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GCAP2 Polyclonal Antibody - Images