

**GPAT2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55182****Specification**

---

**GPAT2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">Q6NUI2</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87835

**GPAT2 Polyclonal Antibody - Additional Information****Gene ID** 150763**Other Names**

Glycerol-3-phosphate acyltransferase 2, mitochondrial, GPAT-2, 2.3.1.15, GPAT2 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=27168" target="\_blank">HGNC:27168</a>)

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**GPAT2 Polyclonal Antibody - Protein Information****Name** GPAT2 ([HGNC:27168](#))**Function**

Transfers an acyl-group from acyl-ACP to the sn-1 position of glycerol-3-phosphate producing a lysophosphatidic acid (LPA), an essential step for the triacylglycerol (TAG) and glycerophospholipids. In vitro also transfers an acyl-group from acyl-ACP to the LPA producing a phosphatidic acid (PA). Prefers arachidonoyl-CoA as the acyl donor. Required for primary processing step during piRNA biosynthesis. Molecular mechanisms by which it promotes piRNA biosynthesis are unclear and do not involve its acyltransferase activity.

**Cellular Location**


Mitochondrion outer membrane {ECO:0000250|UniProtKB:Q14DK4}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q14DK4}

**GPAT2 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### GPAT2 Polyclonal Antibody - Images



Sample: Testis (Mouse) Lysate at 40 ug  
Primary: Anti-GPAT2 (bs-13497R) at 1/300 dilution  
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution  
Predicted band size: 88 kD  
Observed band size: 88 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GPAT2) Polyclonal Antibody, Unconjugated (bs-13497R GPAT2) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.