

HoxC13 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55214**Specification**

HoxC13 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	P31276
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HoxC13
Epitope Specificity	231-330/330
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nuclear
SIMILARITY	Belongs to the Abd-B homeobox family. Contains 1 homeobox DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

The Hox proteins are a family of transcription factors that play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. Hox proteins are involved in controlling axial patterning, leukemias and hereditary malformations. HoxC13 (homeobox C13), also known as HOX3 or HOX3G, is a 330 amino acid protein that contains one homeobox DNA-binding domain and is a member of the Abd-B homeobox family. Localized to the nucleus, HoxC13 functions as a sequence-specific transcription factor that, in conjunction with a variety of other proteins, provides cells with positional identities on their anterior-posterior axis. Via its ability to modify features of the anterior-posterior body axis, HoxC13 is thought to play a role in the development of nails, hair and filiform papilla.

HoxC13 Polyclonal Antibody - Additional Information**Gene ID** 3229**Other Names**

Homeobox protein Hox-C13, Homeobox protein Hox-3G, HOXC13, HOX3G

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
ICC~~N/A
E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HoxC13 Polyclonal Antibody - Protein Information

Name HOXC13

Synonyms HOX3G

Function

Transcription factor which plays a role in hair follicle differentiation. Regulates FOXQ1 expression and that of other hair- specific genes (By similarity).

Cellular Location

Nucleus.

HoxC13 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HoxC13 Polyclonal Antibody - Images