

C19orf80 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55296**Specification**

C19orf80 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	Q6UXH0
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22105

C19orf80 Polyclonal Antibody - Additional Information**Gene ID** 55908**Other Names**

Angiopoietin-like protein 8 {ECO:0000312|HGNC:HGNC:24933}, Betatrophin, Lipasin, Refeeding-induced fat and liver protein, ANGPTL8 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=24933)
target="_blank">HGNC:24933)

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

C19orf80 Polyclonal Antibody - Protein Information**Name** ANGPTL8 ([HGNC:24933](#))**Function**

Hormone that acts as a blood lipid regulator by regulating serum triglyceride levels (PubMed:[22569073](http://www.uniprot.org/citations/22569073), PubMed:[22809513](http://www.uniprot.org/citations/22809513), PubMed:[23150577](http://www.uniprot.org/citations/23150577)). May be involved in the metabolic transition between fasting and refeeding: required to direct fatty acids to adipose tissue for storage in the fed state (By similarity).

Cellular Location

Secreted.

Tissue Location

Predominantly expressed in liver. Also expressed in adipose tissues.

C19orf80 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C19orf80 Polyclonal Antibody - Images