

Centrin 3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55317**Specification**

Centrin 3 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O15182
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	19550

Centrin 3 Polyclonal Antibody - Additional Information**Gene ID** 1070**Other Names**

Centrin-3, CETN3, CEN3

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Centrin 3 Polyclonal Antibody - Protein Information**Name** CETN3**Synonyms** CEN3**Function**

Plays a fundamental role in microtubule-organizing center structure and function.

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus, nucleolus. Nucleus envelope. Nucleus, nuclear pore complex. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Note=Centrosome of interphase and mitotic cells (PubMed:9256449). Localizes to centriole distal lumen (PubMed:26337392). Localization at the nuclear pore complex

requires NUP153 and TPR (PubMed:23591820).

Centrin 3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Centrin 3 Polyclonal Antibody - Images