

CFHL5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55328

Specification

CFHL5 Polyclonal Antibody - Product Information

Application Primary Accession

Host Clonality Calculated MW Physical State Immunogen

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

Buffer

SUBCELLULAR LOCATION SIMILARITY DISEASE IHC-P, IHC-F, IF, ICC, E

Q9BXR6 Rabbit Polyclonal 62 KDa Liquid

KLH conjugated synthetic peptide derived

from human CFHL5

21-120/569

laG

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Secreted.

Contains 9 Sushi (CCP/SCR) domains. Note=Defects in CFHR5 have been found in patients with atypical hemolytic uremic syndrome and may contribute to the disease. Atypical hemolytic uremic syndrome is a complex genetic disease characterized by microangiopathic hemolytic anemia, thrombocytopenia, renal failure and absence of episodes of enterocolitis and diarrhea. In contrast to typical hemolytic uremic syndrome, atypical forms have a poorer prognosis, with higher death rates and frequent progression to end-stage renal disease. Susceptibility to the development of atypical hemolytic uremic syndrome can be conferred by mutations in various components of or regulatory factors in the complement cascade system. Other genes may play a role in modifying the

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Important Note

This gene is a member of a small complement factor H (CFH) gene cluster on chromosome 1. Each member of this gene family contains multiple short consensus repeats (SCRs) typical of regulators of complement activation. The protein encoded by this gene has nine SCRs with the first two repeats having heparin binding properties, a region within repeats 5-7 having heparin binding and

phenotype.



C reactive protein binding properties, and the C-terminal repeats being similar to a complement component 3 b (C3b) binding domain. This protein co-localizes with C3, binds C3b in a dose-dependent manner, and is recruited to tissues damaged by C-reactive protein. Allelic variations in this gene have been associated, but not causally linked, with two different forms of kidney disease: membranoproliferative glomerulonephritis type II (MPGNII) and hemolytic uraemic syndrome (HUS). [provided by RefSeq, Jan 2010]

CFHL5 Polyclonal Antibody - Additional Information

Gene ID 81494

Other Names

Complement factor H-related protein 5, FHR-5, CFHR5, CFHL5, FHR5

Target/Specificity

Expressed by the liver and secreted in plasma.

Dilution

```
<span class ="dilution_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution_IF">IF~~1:50~200</span><br \><span class ="dilution_ICC">ICC~~N/A</span><br \><span class ="dilution_E">E~~N/A</span>
```

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CFHL5 Polyclonal Antibody - Protein Information

Name CFHR5

Synonyms CFHL5, FHR5

Function

Involved in complement regulation. The dimerized forms have avidity for tissue-bound complement fragments and efficiently compete with the physiological complement inhibitor CFH.

Cellular Location

Secreted.


Tissue Location

Expressed by the liver and secreted in plasma.

CFHL5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry





• <u>Immunofluorescence</u>

- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CFHL5 Polyclonal Antibody - Images