

## **CHST5 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55348

### **Specification**

## **CHST5 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC

Primary Accession <u>Q9GZS9</u>

Reactivity
Host
Clonality
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal

Clonality Polycle Calculated MW 46161

# **CHST5 Polyclonal Antibody - Additional Information**

### Gene ID 23563

#### **Other Names**

Carbohydrate sulfotransferase 5, 2.8.2.-, Galactose/N-acetylglucosamine/N-acetylglucosamine 6-O-sulfotransferase 4-alpha, GST4-alpha, Intestinal N-acetylglucosamine-6-O-sulfotransferase, I-GlcNAc6ST, Intestinal GlcNAc-6-sulfotransferase, hIGn6ST, N-acetylglucosamine 6-O-sulfotransferase 3, GlcNAc6ST-3, Gn6st-3, CHST5

### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

### **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# **CHST5 Polyclonal Antibody - Protein Information**

### Name CHST5

### **Function**

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of non-reducing N-acetylglucosamine (GlcNAc) residues and O-linked sugars of mucin-type acceptors. Acts on the non-reducing terminal GlcNAc of short carbohydrate substrates. However, it does not transfer sulfate to longer carbohydrate substrates that have poly-N- acetyllactosamine structures. Has no activity toward keratan. Not involved in generating HEV-expressed ligands for SELL. Its substrate specificity may be influenced by its subcellular location.

### **Cellular Location**

Golgi apparatus membrane; Single-pass type II membrane protein. Note=Golgi membrane, early secretory pathway

# **Tissue Location**



Predominantly expressed in small and large intestines and colon. Weakly expressed in lymphocytes. Not expressed in other tissues. Down-regulated in colonic adenocarcinomas

# **CHST5 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**CHST5 Polyclonal Antibody - Images**