

CHST5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55348

Specification

CHST5 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC

Primary Accession <u>O9GZS9</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 46161

CHST5 Polyclonal Antibody - Additional Information

Gene ID 23563

Other Names

Carbohydrate sulfotransferase 5, 2.8.2.-, Galactose/N-acetylglucosamine/N-acetylglucosamine 6-O-sulfotransferase 4-alpha, GST4-alpha, Intestinal N-acetylglucosamine-6-O-sulfotransferase, I-GlcNAc6ST, Intestinal GlcNAc-6-sulfotransferase, hIGn6ST, N-acetylglucosamine 6-O-sulfotransferase 3, GlcNAc6ST-3, Gn6st-3, CHST5

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> ICC~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CHST5 Polyclonal Antibody - Protein Information

Name CHST5

Function

Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of non-reducing N-acetylglucosamine (GlcNAc) residues and O-linked sugars of mucin-type acceptors. Acts on the non-reducing terminal GlcNAc of short carbohydrate substrates. However, it does not transfer sulfate to longer carbohydrate substrates that have poly-N- acetyllactosamine structures. Has no activity toward keratan. Not involved in generating HEV-expressed ligands for SELL. Its substrate specificity may be influenced by its subcellular location.



Tel: 858.875.1900 Fax: 858.875.199

Cellular Location

Golgi apparatus membrane; Single-pass type II membrane protein. Note=Golgi membrane, early secretory pathway

Tissue Location

Predominantly expressed in small and large intestines and colon. Weakly expressed in lymphocytes. Not expressed in other tissues. Down-regulated in colonic adenocarcinomas

CHST5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

CHST5 Polyclonal Antibody - Images