

# CHTF18 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55350

### Specification

## **CHTF18 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	<u>Q8WVB6</u>
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	107383

#### **CHTF18** Polyclonal Antibody - Additional Information

Gene ID 63922

Other Names Chromosome transmission fidelity protein 18 homolog, hCTF18, CHL12, CHTF18, C16orf41, CTF18

Dilution <span class ="dilution\_WB">WB~~1:1000</span><br \><span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class ="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_ICC">ICC~~N/A</span><br \><span class = "dilution\_ICC">ICC~~N/A

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

## **CHTF18** Polyclonal Antibody - Protein Information

Name CHTF18

Synonyms C16orf41, CTF18

#### Function

Chromosome cohesion factor involved in sister chromatid cohesion and fidelity of chromosome transmission. Component of one of the cell nuclear antigen loader complexes, CTF18-replication factor C (CTF18-RFC), which consists of CTF18, CTF8, DCC1, RFC2, RFC3, RFC4 and RFC5. The CTF18-RFC complex binds to single-stranded and primed DNAs and has weak ATPase activity that is stimulated by the presence of primed DNA, replication protein A (RPA) and by proliferating cell nuclear antigen (PCNA). The CTF18-RFC complex catalyzes the ATP- dependent loading of PCNA onto primed and gapped DNA. Interacts with and stimulates DNA polymerase POLH. During DNA



repair synthesis, involved in loading DNA polymerase POLE at the sites of local damage (PubMed:<a href="http://www.uniprot.org/citations/20227374" target="\_blank">20227374</a>).

**Cellular Location** 

Nucleus. Note=Associates with chromatin during S phase

#### CHTF18 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

**CHTF18 Polyclonal Antibody - Images**