

### **GEP100/IQSEC1 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP55358

#### **Specification**

### **GEP100/IQSEC1 Polyclonal Antibody - Product Information**

Application WB, IHC-P, IHC-F, IF, ICC, E

Primary Accession
Reactivity
Rat, Pig, Dog
Rabbit
Clonality
Calculated MW
Rat, Pig, Dog
Rabbit
Polyclonal
108 KDa

Physical State
Liquid
Immunogen
KLH conjugated synthetic peptide derived

laG

from human GEP100

Epitope Specificity 451-550/963

Isotype
Purity
affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm. Nucleus. At steady state, may

be preferentially cytosolic.

SIMILARITY Belongs to the BRAG family. Contains 1 IQ

domain. Contains 1 PH domain. Contains 1

SEC7 domain.

Important Note

This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

### **Background Descriptions**

The ADP-ribosylation factor (ARF) protein family are structurally and functionally conserved members of the Ras superfamily of regulatory GTP-binding proteins. ARFs influence vesicle trafficking and signal transduction in eukaryotic cells. ARF6 plays a role in protein trafficking near the plasma membrane, including receptor recycling, cell adhesion and cell migration. ARF6 colocalizes with the ARF guanine nucleotide-exchange protein (GEP) BRAG2, also designated GEP100. BRAG2 is ubiquitously expressed as two isoforms, BRAG2a and BRAG2b, which can cycle between the cytoplasm and the nucleus. BRAG2, via its interaction with ARF6, is involved in the regulation of cell adhesion by controlling Integrin ∫1 endocytosis and E-cadherin redistribution. BRAG2 has also been shown to bind directly to Tyr1068/1086-phosphorylated EGFR to activate ARF6, which induces tumor invasion in MCF7 cells. Therefore, BRAG2 may contribute to the metastasis and malignancy of some breast cancer cells.

#### **GEP100/IQSEC1 Polyclonal Antibody - Additional Information**

**Gene ID 9922** 

### **Other Names**

IQ motif and SEC7 domain-containing protein 1, ADP-ribosylation factors guanine



nucleotide-exchange protein 100, ADP-ribosylation factors guanine nucleotide-exchange protein 2, Brefeldin-resistant Arf-GEF 2 protein, BRAG2, IQSEC1 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=29112" target="blank">HGNC:29112</a>)

### **Target/Specificity**

Expressed in brain, ovary, heart, lung, liver, kidney and leukocytes. Moderate expression was also detected in lung, skeletal muscle, placenta, small intestine, pancreas, spleen and testis.

#### **Dilution**

```
<span class ="dilution_WB">WB~~1:1000</span><br \><span class
="dilution_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution_IF">IF~~1:50~200</span><br \><span class ="dilution_ICC">ICC~~N/A</span><br \><span class ="dilution_ICC">ICC~~N/A</span><br \><span class ="dilution_ICC">ICC~~N/A</span>
```

### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### **GEP100/IQSEC1 Polyclonal Antibody - Protein Information**

#### Name IQSEC1 (HGNC:29112)

#### **Function**

Guanine nucleotide exchange factor for ARF1 and ARF6 (PubMed:<a href="http://www.uniprot.org/citations/11226253" target="\_blank">11226253</a>, PubMed:<a href="http://www.uniprot.org/citations/24058294" target="\_blank">24058294</a>). Guanine nucleotide exchange factor activity is enhanced by lipid binding (PubMed:<a href="http://www.uniprot.org/citations/24058294" target="\_blank">24058294</a>). Accelerates GTP binding by ARFs of all three classes. Guanine nucleotide exchange protein for ARF6, mediating internalization of beta-1 integrin (PubMed:<a href="http://www.uniprot.org/citations/16461286" target="\_blank">16461286</a>). Involved in neuronal development (Probable). In neurons, plays a role in the control of vesicle formation by endocytoc cargo. Upon long term depression, interacts with GRIA2 and mediates the activation of ARF6 to internalize synaptic AMPAR receptors (By similarity).

#### **Cellular Location**

Cytoplasm. Nucleus. Postsynaptic density {ECO:0000250|UniProtKB:Q8R0S2}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250|UniProtKB:Q8R0S2}. Note=At steady state, may be preferentially cytosolic

#### **Tissue Location**

Expressed in brain, ovary, heart, lung, liver, kidney and leukocytes. Moderate expression was also detected in lung, skeletal muscle, placenta, small intestine, pancreas, spleen and testis.

#### GEP100/IQSEC1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot








- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **GEP100/IQSEC1 Polyclonal Antibody - Images**



## Sample:

Cerebrum (Mouse) Lysate at 40 ug

Primary: Anti- GEP100/IQSEC1 (bs-13960R) at 1/300 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 108 kD Observed band size: 108 kD